

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
13 June 2002 (13.06.2002)

PCT

(10) International Publication Number
WO 02/46159 A1

(51) International Patent Classification⁷: **C07D 211/34**,
401/12, A61K 31/445

37, 63505 Langenselbold (DE). **AL-OBEIDI, Fahad**;
6798 N. Corte Calabaza, Tuscon, AZ 85704 (US).

(21) International Application Number: PCT/EP01/13874

(22) International Filing Date:
28 November 2001 (28.11.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
00126750.9 6 December 2000 (06.12.2000) EP

(71) Applicant: **AVENTIS PHARMA DEUTSCHLAND GMBH** [DE/DE]; Brüningstraße 50, 65929 Frankfurt (DE).

(72) Inventors: **PEYMAN, Anuschirwan**; Zeilsheimer Strasse 46, 65779 Kelkheim (DE). **WILL, David, William**; Kirchstrasse 21, 65830 Kriftel (DE). **GERLACH, Uwe**; Im Heideck 30, 65795 Hattersheim (DE). **NAZARÉ, Marc**; Schützenstrasse 7, 65817 Eppstein (DE). **ZOLLER, Gerhard**; Höhenstrasse 8, 61137 Schöneck (DE). **NESTLER, Hans-Peter**; Sendelbacher Weg 46, 65779 Kelkheim (DE). **MATTER, Hans**; Feldbergring

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: GUANIDINE AND AMIDINE DERIVATIVES AS FACTOR XA INHIBITORS

(57) Abstract: The present invention relates compounds of the formula (I), in which R₀, Q, X, Q', D, R₁₀ and V have the meanings indicated in the claims. The compounds of the formula I are valuable pharmacologically active compounds. They exhibit a strong antithrombotic effect and are suitable, for example, for the therapy and prophylaxis of cardiovascular disorders like thromboembolic diseases or restenoses. They are reversible inhibitors of the blood clotting enzymes factor Xa(Fxa) and/or factor VIIa(FVIIa), and can in general be applied in conditions in which an undesired activity of factor Xa and/or factor VIIa is present or for the cure or prevention of which an inhibition of factor Xa and/or factor VIIa is intended. The invention furthermore relates to processes for the preparation of compounds of the formula (I), their use, in particular as active ingredients in pharmaceuticals, and pharmaceutical preparations comprising them.




WO 02/46159 A1

Description

GUANIDINE AND AMIDINE DERIVATIVES AS FACTOR Xa INHIBITORS

5

The present invention relates to compounds of the formula I,



10

in which R_0 ; Q; X; Q', D, R_{10} and V have the meanings indicated below. The compounds of the formula I are valuable pharmacologically active compounds. They exhibit a strong antithrombotic effect and are suitable, for example, for the therapy and prophylaxis of cardiovascular disorders like thromboembolic diseases or restenoses. They are reversible inhibitors of the blood clotting enzymes factor Xa (FXa) and/or factor VIIa (FVIIa), and can in general be applied in conditions in which an undesired activity of factor Xa and/or factor VIIa is present or for the cure or prevention of which an inhibition of factor Xa and/or factor VIIa is intended. The invention furthermore relates to processes for the preparation of compounds of the formula I, their use, in particular as active ingredients in pharmaceuticals, and pharmaceutical preparations comprising them.

15

20

25

Normal haemostasis is the result of a complex balance between the processes of clot initiation, formation and clot dissolution. The complex interactions between blood cells, specific plasma proteins and the vascular surface, maintain the fluidity of blood unless injury and blood loss occurs (EP-A-987274). Many significant disease states are related to abnormal haemostasis. For example, local thrombus formation due to rupture of atherosclerotic plaque is a major cause of acute myocardial infarction and unstable angina. Treatment of an occlusive coronary thrombus by either thrombolytic therapy or percutaneous angioplasty may be accompanied by acute thrombolytic reclosure of the affected vessel.

There continues to be a need for safe and effective therapeutic anticoagulants to limit or prevent thrombus formation. It is most desirable to develop agents that inhibit


coagulation without directly inhibiting thrombin but by inhibiting other steps in the coagulation cascade like factor Xa and/or factor VIIa activity. It is now believed that inhibitors of factor Xa carry a lower bleeding risk than thrombin inhibitors (A. E. P. Adang & J. B. M. Rewinkel, *Drugs of the Future* 2000, 25, 369-383).

- 5 Low molecular weight, factor Xa-specific blood clotting inhibitors that are effective but do not cause unwanted side effects have been described, for example, in WO-A-95/29189. However, besides being an effective factor Xa-specific blood clotting inhibitor, it is desirable that such inhibitors also have further advantageous properties, for instance stability in plasma and liver and selectivity versus other serine proteases whose inhibition
10 is not intended, such as thrombin. There is an ongoing need for further low molecular weight factor Xa specific blood clotting inhibitors which are effective and have the above advantages as well.

- Specific inhibition of the factor VIIa/tissue factor catalytic complex using monoclonal
15 antibodies (WO-A-92/06711) or a protein such as chloromethyl ketone inactivated factor VIIa (WO-A-96/12800, WO-A-97/47651) is an extremely effective means of controlling thrombus formation caused by acute arterial injury or the thrombotic complications related to bacterial septicemia. There is also experimental evidence suggesting that inhibition of factor VIIa/tissue factor activity inhibits restenosis following balloon
20 angioplasty. Bleeding studies have been conducted in baboons and indicate that inhibition of the factor VIIa/tissue factor complex has the widest safety window with respect to therapeutic effectiveness and bleeding risk of any anticoagulant approach tested including thrombin, platelet and factor Xa inhibition. Certain inhibitors of factor VIIa have already been described. EP-A-987274, for example discloses compounds
25 containing a tripeptide unit which inhibit factor VIIa. However, the property profile of these compounds is still not ideal, and there is an ongoing need for further low molecular weight factor VIIa inhibitory blood clotting inhibitors.

- The present invention satisfies the above needs by providing novel compounds of the
30 formula I which exhibit factor Xa and/or factor VIIa inhibitory activity and are favorable agents for inhibiting unwanted blood clotting and thrombus formation.

Thus, the present invention relates to compounds of the formula I,



wherein

- R_0 is
1. phenyl, wherein phenyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R^2 or
 2. a mono- or bicyclic 5- to 10-membered heteroaryl containing one or two nitrogen atoms as ring heteroatoms, wherein heteroaryl is unsubstituted or mono-, di- or trisubstituted independently of one another by R^2 ,

- R^2 is
1. $-\text{NO}_2$,
 2. halogen,
 3. $-\text{CN}$,
 4. $-\text{OH}$,
 5. $-\text{NH}_2$,
 6. $(\text{C}_1\text{-C}_8)\text{-alkyloxy-}$, wherein alkyloxy is unsubstituted or mono-, di- or trisubstituted independently of one another by halogen, amino group, hydroxy group or methoxy group, or
 7. $-(\text{C}_1\text{-C}_8)\text{-alkyl}$, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by halogen, amino group, hydroxy group or methoxy group,

Q and Q' are independently of one another identical or different and are a direct bond, $-\text{O}-$, $-\text{S}-$, $-\text{NR}^{10}-$, $-\text{C}(\text{O})\text{NR}^{10}-$, $-\text{NR}^{10}\text{C}(\text{O})-$, $-\text{S}(\text{O})-$, $-\text{SO}_2-$, $-\text{NR}^{10}\text{-SO}_2-$, $-\text{SO}_2\text{-NR}^{10}-$ oder $-\text{C}(\text{O})-$;


R^{10} is hydrogen atom or $(\text{C}_1\text{-C}_4)\text{-alkyl-}$,

- X is
1. a direct bond,
 2. $(\text{C}_1\text{-C}_6)\text{-alkylen}$, wherein alkylen is unsubstituted or mono-, di- or trisubstituted independently of one another by halogen, amino group or hydroxy group,
 3. $(\text{C}_3\text{-C}_6)\text{-cycloalkylen}$, wherein cycloalkylen is unsubstituted or mono-, di- or trisubstituted independently of one another by halogen, amino group or hydroxy group,

provided that at least one of Q , X and Q' is not a direct bond,

D is an atom out of the group carbon, oxygen, sulfur and nitrogen,

the substructure of formula III



(III)

- is
1. a mono- or bicyclic 5- to 10-membered carbocyclic aryl group, wherein said 5- to 10-membered carbocyclic aryl group is unsubstituted or mono-, di- or trisubstituted independently of one another by R^1 ,
 2. phenyl, wherein phenyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R^1 ,
 3. a mono- or bicyclic 5-to 10-membered heterocyclic group (Het), containing one or more heteroatoms as ring heteroatoms, such as nitrogen, sulfur or oxygen, wherein said Het group is unsubstituted or mono-, di- or trisubstituted independently of one another by R^1 , or
 4. pyridyl, wherein pyridyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R^1 ,

- R^1 is
1. halogen,
 2. $-NO_2$,
 3. $-CN$,
 4. $R^{11}R^{12}N-$, wherein $R^{11}R^{12}$ independently of one another are hydrogen atom, (C_1-C_4) -alkyl- or (C_1-C_6) -acyl-,
 5. (C_1-C_8) -alkylamino-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R^{13} ,
 6. $-OH$,
 7. $-SO_2-NH_2$,
 8. (C_1-C_8) -alkyloxy-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R^{13} ,
 9. (C_6-C_{14}) -aryl, wherein aryl is unsubstituted or mono-, di- or trisubstituted independently of one another by R^{13} ,
 10. (C_1-C_8) -alkyl-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R^{13} ,
 11. hydroxycarbonyl- (C_1-C_8) -alkylureido-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R^{13} ,







12. (C₁-C₈)-alkyloxycarbonyl-(C₁-C₈)-alkylureido-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³,
13. (C₁-C₈)-alkylsulfonyl-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³, or
14. -C(O)-NR¹⁴R¹⁵, wherein R¹⁴R¹⁵ independently of one another are hydrogen atom or (C₁-C₄)-alkyl-, or

two R¹ residues bonded to adjacent ring carbon atoms together with the carbon atoms to which they are bonded form an aromatic ring condensed to the ring depicted in formula I, where the ring formed by the two R¹ residues is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³, or

R¹¹ and R¹² together with the nitrogen atom to which they are bonded form a saturated or unsaturated 5- to 6-membered monocyclic heterocyclic ring which in addition to the nitrogen atom carrying R¹¹ and R¹² can contain one or two identical or different ring heteroatoms chosen from oxygen, sulfur and nitrogen, and in which one or two of the ring carbon atoms can be substituted by oxo to form -C(O)-residue(s),

- R¹³ is
1. halogen,
 2. -NO₂,
 3. -CN,
 4. -OH,
 5. (C₁-C₈)-alkyl-,
 6. (C₁-C₈)-alkyloxy-,
 7. -CF₃ or
 8. -NH₂,

V is a residue of the formulae IIa, IIb, IIc, IId, IIe or IIf,



wherein

L is a direct bond or (C₁-C₃)-alkylene, wherein alkylene is unsubstituted or mono-, di- or trisubstituted independently of one another by A,

- 5 A is
1. hydrogen atom,
 2. -C(O)-OH,
 3. -C(O)-O-(C₁-C₄)-alkyl, wherein alkyl is unsubstituted or mono-, di- or tri-substituted independently of one another by -OH, -NH₂ or -(C₁-C₄)-alkoxy,
 4. -C(O)-NR⁴R⁵,
 - 10 5. (C₁-C₄)-alkyl-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by -OH, -NH₂ or -(C₁-C₄)-alkoxy,
 6. -SO₂-NH₂ or
 7. -SO₂-CH₃,

U is -NH₂, (C₁-C₄)-alkyl-, -NH-C(O)-O-(C₁-C₄)-alkyl or -NH-C(O)-O-(C₁-C₄)-alkyl-aryl,

15 M is hydrogen atom, (C₁-C₃)-alkyl- or -OH,

R⁴ and R⁵ are independently of one another identical or different and are

1. hydrogen atom,
2. (C₁-C₁₂)-alkyl-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³ as defined above,
- 20 3. (C₆-C₁₄)-aryl-(C₁-C₄)-alkyl-, wherein alkyl and aryl are unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³ as defined above,

4. (C₆-C₁₄)-aryl-, wherein aryl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³ as defined above,
 5. Het-, wherein Het- is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³ as defined above, or
 - 5 6. Het-(C₁-C₄)-alkyl-, wherein alkyl and Het- are unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³ as defined above, or
- R⁴ and R⁵ together with the nitrogen atom to which they are bonded form a saturated 3- to 8-membered monocyclic heterocyclic ring which in addition to the nitrogen atom carrying R⁴ and R⁵ can contain one or two identical or different ring heteroatoms
- 10 chosen from oxygen, sulfur and nitrogen;
- in all its stereoisomeric forms and mixtures thereof in any ratio, and its physiologically tolerable salts.

Preferred are compounds of the formula I, wherein

- 15 R₀ is phenyl, wherein phenyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R², or
- pyridyl, wherein pyridyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R²,
- R² is
1. -NO₂,
 - 20 2. halogen,
 3. -CN,
 4. -OH,
 5. -NH₂,
 6. (C₁-C₄)-alkyloxy-, wherein alkyloxy is unsubstituted or mono-, di- or
 - 25 trisubstituted independently of one another by halogen, amino group, hydroxy group or methoxy group, , or
 7. -(C₁-C₄)-alkyl, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by halogen, amino group, hydroxy group or methoxy group,
- 30 Q, Q' , X, R¹, R¹¹ and R¹² are as defined above,
- D is an atom out of the group carbon and nitrogen,
- the substructure of formula III is
1. phenyl, wherein phenyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹, or

2. pyridyl, wherein pyridyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R^1 ,

- R^{13} is
1. halogen,
 2. $-NO_2$,
 3. $-CN$,
 4. $-OH$,
 5. (C_1-C_4) -alkyl-,
 6. (C_1-C_4) -alkyloxy-,
 7. $-CF_3$ or
 8. $-NH_2$,

R_{10} is hydrogen atom or methyl,

V is a fragment of the formula IIa, IIb, IIc, IId, IIe or II f as defined above, wherein

L, U, M, R^4 and R^5 are as defined above, and

- A is
1. hydrogen atom,
 2. $-C(O)-OH$,
 3. $-C(O)-O-(C_1-C_4)$ -alkyl, wherein alkyl is unsubstituted or mono-, di- or tri-substituted independently of one another by $-OH$, $-NH_2$ or $-(C_1-C_4)$ -alkoxy,
 4. $-C(O)-NR^4R^5$ or
 5. (C_1-C_4) -alkyl-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by $-OH$, $-NH_2$ or $-(C_1-C_4)$ -alkoxy.

More preferred are compounds of the formula I, wherein

R^0 is phenyl, wherein phenyl is mono-, di- or trisubstituted independently of one another by R^2 , or

pyridyl, wherein pyridyl is mono-, di- or trisubstituted independently of one another by R^2 ,

- R^2 is
1. $-NH_2$,
 2. halogen,
 3. $-CN$,
 4. $-OH$,
 5. (C_1-C_4) -alkyloxy-, wherein alkyloxy is unsubstituted or substituted by an amino group, or
 6. $-(C_1-C_4)$ -alkyl, wherein alkyl is unsubstituted or substituted by an amino group,

Q and Q' are independently of one another identical or different and are

a direct bond, -O-, -C(O)NR¹⁰-, -NR¹⁰C(O)-; -NR¹⁰-SO₂-; or -SO₂-NR¹⁰-;

X is 1. a direct bond or

2. (C₁-C₄)-alkylen, wherein alkylen is unsubstituted or mono-, di- or tri-substituted independently of one another by halogen, amino group or hydroxy group,

D is an atom out of the group carbon and nitrogen,

the substructure of formula III is

phenyl or pyridyl, wherein phenyl and pyridyl are unsubstituted or mono-,

di- or trisubstituted independently of one another by R¹,

R¹ is 1. halogen,

2. -NO₂,

3. -CN,

4. -NH₂,

5. (C₁-C₄)-alkylamino-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³,

6. -OH,

7. -SO₂-NH₂,

8. (C₁-C₄)-alkyloxy-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³,

9. (C₆-C₁₄)-aryl, wherein aryl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³,

10. (C₁-C₄)-alkyl-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³,

11. (C₁-C₄)-alkylsulfonyl-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³,

12. -C(O)-NR¹⁴R¹⁵, wherein R¹⁴R¹⁵ independently of one another are hydrogen atom or (C₁-C₄)-alkyl-,

13. R¹¹R¹²N-, wherein R¹¹ and R¹² are as defined above, or

14. -NR⁴R⁵,

R¹³ is 1. halogen,

2. -NO₂,

3. -CN,

4. -OH,

5. (C₁-C₄)-alkyl-,
6. (C₁-C₄)-alkyloxy-,
7. -CF₃ or
8. -NH₂,

5 R₁₀ is hydrogen atom or methyl,

V is a fragment of the formula IIa, IIb, IIc, IId, IIe or II f as defined above, wherein

L is a direct bond or (C₁-C₃)-alkylen-,

A is hydrogen atom, -C(O)-OH, -C(O)-O-(C₁-C₄)-alkyl, -C(O)-NR⁴R⁵ or (C₁-C₄)-alkyl,

U is -NH₂, methyl, -NH-C(O)-O-(C₁-C₄)-alkyl or -NH-C(O)-O-(CH₂)-phenyl,

10 M is hydrogen atom, (C₁-C₃)-alkyl- or -OH, and

R⁴ and R⁵ are independently of one another hydrogen atom or (C₁-C₄)-alkyl-.

Even more preferred are the compounds of the formula I, wherein

R₀ is phenyl or pyridyl, wherein phenyl and pyridyl independently from one another are

15 mono-, di- or trisubstituted independently of one another by R²,

R² is 1. halogen,

2. -CN,

3. (C₁-C₄)-alkyloxy-, wherein alkyloxy is unsubstituted or substituted by halogen or an amino group, or

20 4. -(C₁-C₄)-alkyl, wherein alkyl is unsubstituted or substituted by an amino group or halogen,

Q and Q' are independently of one another identical or different and are

a direct bond, -O-, -C(O)NR¹⁰-, -NR¹⁰C(O)-; -NR¹⁰-SO₂-; or -SO₂-NR¹⁰-;

X is -(C₁-C₃)-alkylen-, wherein alkylen is unsubstituted or mono-, di- or trisubstituted

25 independently of one another by halogen, amino group or hydroxy group,

D is the atom carbon,

the substructure of formula III is

phenyl, wherein phenyl is unsubstituted or mono-, di- or trisubstituted

independently of one another by R¹,

30 R¹ is 1. halogen,

2. -NO₂,

3. -CN,

4. -NH₂,

5. (C₁-C₄)-alkylamino-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³,
6. -OH,
7. -SO₂-NH₂,
- 5 8. (C₁-C₄)-alkyloxy-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³,
9. (C₆-C₁₄)-aryl, wherein aryl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³,
- 10 10. (C₁-C₄)-alkyl-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³,
11. (C₁-C₄)-alkylsulfonyl-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³,
12. -C(O)-NR¹⁴R¹⁵, wherein R¹⁴R¹⁵ independently of one another are hydrogen atom or (C₁-C₄)-alkyl-,
- 15 13. R¹¹R¹²N-, wherein R¹¹ and R¹² are as defined above, or
14. -NR⁴R⁵, wherein R⁴ and R⁵ are independently of one another hydrogen atom or methyl

- R¹³ is 1. halogen,
2. -CF₃,
- 20 3. -NH₂,
4. -OH,
5. (C₁-C₄)-alkyl- or
6. (C₁-C₄)-alkyloxy-,

R₁₀ is hydrogen atom, and

25 V is a fragment of the formula IIa, IIb, IIc or IId as defined above, wherein

L is a direct bond or (C₁-C₂)-alkylen-,

A is hydrogen atom, -C(O)-OH, -C(O)-O-(C₁-C₄)-alkyl, -C(O)-NR⁴R⁵ or (C₁-C₄)-alkyl,

U is -NH₂, methyl, -NH-C(O)-O-(C₁-C₄)-alkyl or -NH-C(O)-O-(CH₂)-phenyl,

M is hydrogen atom or (C₁-C₃)-alkyl-.

30

Further preferred are compounds of the formula I, wherein

R₀ is phenyl, wherein phenyl is mono-, di- or trisubstituted independently of one another by R²,

R² is 1. halogen,

2. (C₁-C₄)-alkyloxy-, wherein alkyloxy is unsubstituted or substituted by halogen or an amino group, or
3. -(C₁-C₄)-alkyl, wherein alkyl is unsubstituted or substituted by an amino group or halogen,

5 Q and Q' are independently of one another identical or different and are a direct bond, -O-, -C(O)NR¹⁰-, -NR¹⁰C(O)-; -NR¹⁰-SO₂-; or -SO₂-NR¹⁰-;

X is -(C₁-C₃)-alkylen-,

D is the atom carbon,

the substructure of formula III is

10 phenyl, wherein phenyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹,

R¹ is 1. halogen,

2. -NO₂,

3. -CN,

15 4. -NH₂,

5. (C₁-C₄)-alkylamino-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³,

6. -OH,

7. -SO₂-NH₂,

20 8. (C₁-C₄)-alkyloxy-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³,

9. (C₁-C₄)-alkyl-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³,

25 10. (C₁-C₄)-alkylsulfonyl-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³,

11. -C(O)-NR¹⁴R¹⁵, wherein R¹⁴R¹⁵ independently of one another are hydrogen atom or (C₁-C₂)-alkyl-,

12. R¹¹R¹²N-, wherein R¹¹ and R¹² are as defined above, or

13. -NR⁴R⁵,

30 R¹³ is 1. halogen,

2. -CF₃,

3. -NH₂,

4. -OH,

5. (C₁-C₄)-alkyl- or

6. (C₁-C₄)-alkyloxy-,

R₁₀ is hydrogen atom, and

V is a fragment of the formula IIa, IIb, IIc or IId as defined above, wherein

L is a direct bond or (C₁-C₂)-alkylen-,

5 A is hydrogen atom, -C(O)-OH, -C(O)-O-(C₁-C₄)-alkyl, -C(O)-NR⁴R⁵ or -(C₁-C₄)-alkyl,

U is -NH₂, methyl, -NH-C(O)-O-(C₁-C₄)-alkyl or -NH-C(O)-O-(CH₂)-phenyl,

M is hydrogen atom or methyl, and

R⁴ and R⁵ are independently of one another hydrogen atom or methyl.

10 Particularly preferred are compounds of the formula I, wherein

R₀ is phenyl, wherein phenyl is disubstituted independently of one another by R²,

R² is 1. halogen,

2. (C₁-C₂)-alkyloxy-, wherein alkyloxy is unsubstituted or substituted by an amino group, or

15 3. -(C₁-C₄)-alkyl, wherein alkyl is unsubstituted or substituted by an amino group,

Q and Q' are independently of one another identical or different and are

a direct bond or -O-,

X is -CH₂-CH₂-,

20 D is the atom carbon,

the substructure of formula III is

phenyl, wherein phenyl is unsubstituted or mono-, di- or trisubstituted

independently of one another by R¹,

R¹ is 1. halogen,

25 2. -OH,

3. -NH₂,

4. -C(O)-NR¹⁴R¹⁵, wherein R¹⁴R¹⁵ independently of one another are hydrogen atom or (C₁-C₂)-alkyl-,

5. (C₁-C₃)-alkyloxy-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³, or

30 6. (C₁-C₃)-alkyl-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³,

R¹³ is fluorine or chlorine,

R₁₀ is hydrogen atom, and

V is a fragment of the formula IIa, IIb, IIc or IId as defined above, wherein

L is a direct bond or (C₁-C₂)-alkylen-,

A is hydrogen atom, -C(O)-OH, -C(O)-O-(C₁-C₄)-alkyl, -C(O)-NR⁴R⁵ or -(C₁-C₄)-alkyl,

U is -NH₂, methyl, -NH-C(O)-O-(C₁-C₄)-alkyl or -NH-C(O)-O-(CH₂)-phenyl,

5 M is hydrogen atom, and

R⁴ and R⁵ are independently of one another hydrogen atom or methyl.

In general, the meaning of any group, residue, heteroatom, number etc. which can occur more than once in the compounds of the formula I, is independent of the meaning of this
10 group, residue, heteroatom, number etc. in any other occurrence. All groups, residues, heteroatoms, numbers etc. which can occur more than once in the compounds of the formula I can be identical or different.

As used herein, the term alkyl is to be understood in the broadest sense to mean hydrocarbon residues which can be linear, i. e. straight-chain, or branched and which
15 can be acyclic or cyclic residues or comprise any combination of acyclic and cyclic subunits. Further, the term alkyl as used herein expressly includes saturated groups as well as unsaturated groups which latter groups contain one or more, for example one, two or three, double bonds and/or triple bonds, provided that the double bonds are not located within a cyclic alkyl group in such a manner that an aromatic system results. All
20 these statements also apply if an alkyl group occurs as a substituent on another residue, for example in an alkyloxy residue, an alkyloxycarbonyl residue or an arylalkyl residue. Examples of alkyl residues containing 1, 2, 3, 4, 5, 6, 7 or 8 carbon atoms are methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl or octyl, the n-isomers of all these residues, isopropyl, isobutyl, 1-methylbutyl, isopentyl, neopentyl, 2,2-dimethylbutyl, 2-methylpentyl,
25 3-methylpentyl, isohexyl, sec-butyl, tert-butyl, tert-pentyl, sec-butyl, tert-butyl or tert-pentyl.

Unsaturated alkyl residues are, for example, alkenyl residues such as vinyl, 1-propenyl, 2-propenyl (= allyl), 2-butenyl, 3-butenyl, 2-methyl-2-butenyl, 3-methyl-2-butenyl, 5-
30 hexenyl or 1,3-pentadienyl, or alkynyl residues such as ethynyl, 1-propynyl, 2-propynyl (= propargyl) or 2-butyne. Alkyl residues can also be unsaturated when they are substituted.

Examples of cyclic alkyl residues are cycloalkyl residues containing 3, 4, 5 or 6 ring carbon atoms like cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl, which can also be substituted and/or unsaturated. Unsaturated cyclic alkyl groups and unsaturated cycloalkyl groups like, for example, cyclopentenyl or cyclohexenyl can be bonded via any carbon atom.

Of course, a cyclic alkyl group has to contain at least three carbon atoms, and an unsaturated alkyl group has to contain at least two carbon atoms. Thus, a group like (C₁-C₈)-alkyl is to be understood as comprising, among others, saturated acyclic (C₁-C₈)-alkyl, (C₃-C₆)-cycloalkyl, and unsaturated (C₂-C₈)-alkyl like (C₂-C₈)-alkenyl or (C₂-C₈)-alkynyl. Similarly, a group like (C₁-C₄)-alkyl is to be understood as comprising, among others, saturated acyclic (C₁-C₄)-alkyl, and unsaturated (C₂-C₄)-alkyl like (C₂-C₄)-alkenyl or (C₂-C₄)-alkynyl.

Unless stated otherwise, the term alkyl preferably comprises acyclic saturated hydrocarbon residues which have from one to six carbon atoms and which can be linear or branched. A particular group of saturated acyclic alkyl residues is formed by (C₁-C₄)-alkyl residues like methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl and tert-butyl.

Unless stated otherwise, and irrespective of any specific substituents bonded to alkyl groups which are indicated in the definition of the compounds of the formula I, alkyl groups can in general be unsubstituted or substituted by one or more, for example one, two or three, identical or different substituents. Any kind of substituents present in substituted alkyl residues can be present in any desired position provided that the substitution does not lead to an unstable molecule. Examples of substituted alkyl residues are alkyl residues in which one or more, for example 1, 2 or 3, hydrogen atoms are replaced with halogen atoms, in particular fluorine atoms.

The term mono- or bicyclic 5- to 10-membered carbocyclic aryl group refers to for example phenyl or naphthyl.

The term mono- or bicyclic 5- to 10-membered heteroaryl containing one or two nitrogen atoms as ring heteroatoms refers to (C₅-C₁₀)-aryl in which one or more of the 5 to 10 ring carbon atoms are replaced by heteroatoms such as nitrogen, oxygen or sulfur. Examples are pyridyl; such as 2-pyridyl, 3-pyridyl or 4-pyridyl; pyrrolyl; such as 2-pyrrolyl and 3-

pyrrolyl; furyl; such as 2-furyl and 3-furyl; thienyl; such as 2-thienyl and 3-thienyl; imidazolyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, tetrazolyl, pyridazinyl, pyrazinyl, pyrimidinyl, indolyl, isoindolyl, indazolyl, phthalazinyl, quinolyl, isoquinolyl or quinoxalinyl.

5

The term R¹¹ and R¹² together with the nitrogen atom to which they are bonded form a saturated or unsaturated 5- to 6-membered monocyclic heterocyclic ring refers to pyrrol, piperidin, pyrrolidine, morpholine, piperazine, pyridine, pyrimidine, imidazole or thiomorpholine.

10

The term aryl refers to a monocyclic or polycyclic hydrocarbon residue in which at least one carbocyclic ring is present that has a conjugated pi electron system. In a (C₆-C₁₄)-aryl residue from 6 to 14 ring carbon atoms are present. Examples of (C₆-C₁₄)-aryl residues are phenyl, naphthyl, biphenyl, fluorenyl or anthracenyl. Unless stated

15

otherwise, and irrespective of any specific substituents bonded to aryl groups which are indicated in the definition of the compounds of the formula I, aryl residues, for example phenyl, naphthyl or fluorenyl, can in general be unsubstituted or substituted by one or more, for example one, two or three, identical or different substituents. Aryl residues can be bonded via any desired position, and in substituted aryl residues the substituents can

20

Unless stated otherwise, and irrespective of any specific substituents bonded to aryl groups which are indicated in the definition of the compounds of the formula I, substituents that can be present in substituted aryl groups are, for example, (C₁-C₈)-alkyl, in particular (C₁-C₄)-alkyl, such as methyl, ethyl or tert-butyl, hydroxy, (C₁-C₈)-alkyloxy, in particular (C₁-C₄)-alkyloxy, such as methoxy, ethoxy or tert-butoxy, methylenedioxy, ethylenedioxy, F, Cl, Br, I, cyano, nitro, trifluoromethyl, trifluoromethoxy, hydroxymethyl, formyl, acetyl, amino, mono- or di-(C₁-C₄)-alkylamino, ((C₁-C₄)-alkyl)carbonylamino like acetylamino, hydroxycarbonyl, ((C₁-C₄)-alkyloxy)carbonyl, carbamoyl, benzyl optionally substituted in the phenyl group, optionally substituted phenyl, optionally substituted phenoxy or benzyloxy optionally substituted in the phenyl group. A substituted aryl group which is present in a specific position of the compounds of formula I can independently of other aryl groups be substituted by substituents selected from any desired subgroup of the substituents listed before and/or in the specific definition of that group. For example,

30

a substituted aryl group may be substituted by one or more identical or different substituents chosen from (C₁-C₄)-alkyl, hydroxy, (C₁-C₄)-alkyloxy, F, Cl, Br, I, cyano, nitro, trifluoromethyl, amino, phenyl, benzyl, phenoxy and benzyloxy. In general, preferably not more than two nitro groups are present in the compounds of the formula I.

5

In monosubstituted phenyl residues the substituent can be located in the 2-position, the 3-position or the 4-position, with the 3-position and the 4-position being preferred. If a phenyl group carries two substituents, they can be located in 2,3-position, 2,4-position, 2,5-position, 2,6-position, 3,4-position or 3,5-position. In phenyl residues carrying three substituents the substituents can be located in 2,3,4-position, 2,3,5-position, 2,3,6-position, 2,4,5-position, 2,4,6-position, or 3,4,5-position. Naphthyl residues can be 1-naphthyl and 2-naphthyl. In substituted naphthyl residues the substituents can be located in any positions, for example in monosubstituted 1-naphthyl residues in the 2-, 3-, 4-, 5-, 6-, 7-, or 8-position and in monosubstituted 2-naphthyl residues in the 1-, 3-, 4-, 5-, 6-, 7-, or 8-position. Biphenyl residues can be 2-biphenyl, 3-biphenyl and 4-biphenyl. Fluorenyl residues can be 1-, 2-, 3-, 4- or 9-fluorenyl. In monosubstituted fluorenyl residues bonded via the 9-position the substituent is preferably present in the 1-, 2-, 3- or 4-position.

20 The group Het comprises groups containing 5, 6, 7, 8, 9 or 10 ring atoms in the parent monocyclic or bicyclic heterocyclic ring system. In monocyclic groups Het the heterocyclic ring preferably is a 5-membered, 6-membered or 7-membered ring, particularly preferably a 5-membered or 6-membered ring. In bicyclic groups Het preferably two fused rings are present one of which is a 5-membered ring or 6-membered heterocyclic ring and the other of which is a 5-membered or 6-membered heterocyclic or carbocyclic ring, i. e. a bicyclic ring Het preferably contains 8, 9 or 10 ring atoms, particularly preferably 9 or 10 ring atoms.

30 Het comprises saturated heterocyclic ring systems which do not contain any double bonds within the rings, as well as mono-unsaturated and poly-unsaturated heterocyclic ring systems which contain one or more, for example one, two, three, four or five, double bonds within the rings provided that the resulting system is stable. Unsaturated rings may be non-aromatic or aromatic, i. e. double bonds within the rings in the group Het may be arranged in such a manner that a conjugated pi electron system results.

Aromatic rings in a group Het may be 5-membered or 6-membered rings, i. e. aromatic groups in a group Het contain 5 to 10 ring atoms. Aromatic rings in a group Het thus comprise 5-membered and 6-membered monocyclic heterocycles and bicyclic heterocycles composed of two 5-membered rings, one 5-membered ring and one 6-membered ring, or two 6-membered rings. In bicyclic aromatic groups in a group Het one or both rings may contain heteroatoms. Aromatic groups Het may also be referred to by the customary term heteroaryl for which all the definitions and explanations above and below relating to Het correspondingly apply.

- 10 Unless stated otherwise, in the groups Het and any other heterocyclic groups preferably 1, 2, 3 or 4 identical or different ring heteroatoms chosen from nitrogen, oxygen and sulfur are present. Particularly preferably in these groups one or two identical or different heteroatoms chosen from nitrogen, oxygen and sulfur are present. The ring heteroatoms can be present in any desired number and in any position with respect to each other
- 15 provided that the resulting heterocyclic system is known in the art and is stable and suitable as a subgroup in a drug substance. Examples of parent structures of heterocycles from which the group Het can be derived are aziridine, oxirane, azetidine, pyrrole, furan, thiophene, dioxole, imidazole, pyrazole, oxazole, isoxazole, thiazole, isothiazole, 1,2,3-triazole, 1,2,4-triazole, tetrazole, pyridine, pyran, thiopyran, pyridazine,
- 20 pyrimidine, pyrazine, 1,2-oxazine, 1,3-oxazine, 1,4-oxazine, 1,2-thiazine, 1,3-thiazine, 1,4-thiazine, 1,2,3-triazine, 1,2,4-triazine, 1,3,5-triazine, azepine, 1,2-diazepine, 1,3-diazepine, 1,4-diazepine, indole, isoindole, benzofuran, benzothiophene, 1,3-benzodioxole, indazole, benzimidazole, benzoxazole, benzothiazole, quinoline, isoquinoline, chromane, isochromane, cinnoline, quinazoline, quinoxaline, phthalazine,
- 25 pyridoimidazoles, pyridopyridines, pyridopyrimidines, purine, pteridine etc. as well as ring systems which result from the listed heterocycles by fusion (or condensation) of a carbocyclic ring, for example benzo-fused, cyclopenta-fused, cyclohexa-fused or cyclohepta-fused derivatives of these heterocycles.
- 30 The fact that many of the before-listed names of heterocycles are the chemical names of unsaturated or aromatic ring systems does not imply that the groups Het could only be derived from the respective unsaturated ring system. The names here only serve to describe the ring system with respect to ring size and the number of the heteroatoms and their relative positions. As explained above, the group Het can be saturated or partially

unsaturated or aromatic, and can thus be derived not only from the before-listed heterocycles themselves but also from all their partially or completely hydrogenated analogues and also from their more highly unsaturated analogues if applicable. As

5 heterocycles from which the groups Het may be derived the following may be mentioned: pyrroline, pyrrolidine, tetrahydrofuran, tetrahydrothiophene, dihydropyridine, tetrahydropyridine, piperidine, 1,3-dioxolane, 2-imidazoline, imidazolidine, 4,5-dihydro-1,3-oxazol, 1,3-oxazolidine, 4,5-dihydro-1,3-thiazole, 1,3-thiazolidine, perhydro-1,4-dioxane, piperazine, perhydro-1,4-oxazine (= morpholine), perhydro-1,4-thiazine (= 10 thiomorpholine), perhydroazepine, indoline, isoindoline, 1,2,3,4-tetrahydroquinoline, 1,2,3,4-tetrahydroisoquinoline, etc.

The residue Het may be bonded via any ring carbon atom, and in the case of nitrogen heterocycles via any suitable ring nitrogen atom. Thus, for example, a pyrrolyl residue 15 can be 1-pyrrolyl, 2-pyrrolyl or 3-pyrrolyl, a pyrrolidinyl residue can be pyrrolidin-1-yl (= pyrrolidino), pyrrolidin-2-yl or pyrrolidin-3-yl, a pyridinyl residue can be pyridin-2-yl, pyridin-3-yl or pyridin-4-yl, a piperidinyl residue can be piperidin-1-yl (= piperidino), piperidin-2-yl, piperidin-3-yl or piperidin-4-yl. Furyl can be 2-furyl or 3-furyl, thienyl can be 2-thienyl or 3-thienyl, imidazolyl can be imidazol-1-yl, imidazol-2-yl, imidazol-4-yl or 20 imidazol-5-yl, 1,3-oxazolyl can be 1,3-oxazol-2-yl, 1,3-oxazol-4-yl or 1,3-oxazol-5-yl, 1,3-thiazolyl can be 1,3-thiazol-2-yl, 1,3-thiazol-4-yl or 1,3-thiazol-5-yl, pyrimidinyl can be pyrimidin-2-yl, pyrimidin-4-yl (= 6-pyrimidinyl) or 5-pyrimidinyl, piperazinyl can be piperazin-1-yl (= piperazin-4-yl = piperazino) or piperazin-2-yl. Indolyl can be indol-1-yl, indol-2-yl, indol-3-yl, indol-4-yl, indol-5-yl, indol-6-yl or indol-7-yl. Similarly 25 benzimidazolyl, benzoxazolyl and benzothiazol residues can be bonded via the 2-position and via any of the positions 4, 5, 6, and 7. Quinoliny can be quinolin-2-yl, quinolin-3-yl, quinolin-4-yl, quinolin-5-yl, quinolin-6-yl, quinolin-7-yl or quinolin-8-yl, isoquinoliny can be isoquinol-1-yl, isoquinolin-3-yl, isoquinolin-4-yl, isoquinolin-5-yl, isoquinolin-6-yl, isoquinolin-7-yl or isoquinolin-8-yl. In addition to being bonded via any of 30 the positions indicated for quinoliny and isoquinoliny, 1,2,3,4-tetrahydroquinoliny and 1,2,3,4-tetrahydroisoquinoliny can also be bonded via the nitrogen atoms in 1-position and 2-position, respectively.

- Unless stated otherwise, and irrespective of any specific substituents bonded to groups Het or any other heterocyclic groups which are indicated in the definition of the compounds of the formula I, the group Het can be unsubstituted or substituted on ring carbon atoms with one or more, for example one, two, three, four or five, identical or different substituents like (C₁-C₈)-alkyl, in particular (C₁-C₄)-alkyl, (C₁-C₈)-alkyloxy, in particular (C₁-C₄)-alkyloxy, (C₁-C₄)-alkylthio, halogen, nitro, amino, ((C₁-C₄)-alkyl)carbonylamino like acetylamino, trifluoromethyl, trifluoromethoxy, hydroxy, oxo, hydroxy-(C₁-C₄)-alkyl such as, for example, hydroxymethyl or 1-hydroxyethyl or 2-hydroxyethyl, methylenedioxy, ethylenedioxy, formyl, acetyl, cyano, aminosulfonyl, methylsulfonyl, hydroxycarbonyl, aminocarbonyl, (C₁-C₄)-alkyloxycarbonyl, optionally substituted phenyl, optionally substituted phenoxy, benzyl optionally substituted in the phenyl group, benzyloxy optionally substituted in the phenyl group, etc. The substituents can be present in any desired position provided that a stable molecule results. Of course an oxo group cannot be present in an aromatic ring. Each suitable ring nitrogen atom in a group Het can independently of each other be unsubstituted, i. e. carry a hydrogen atom, or can be substituted, i. e. carry a substituent like (C₁-C₈)-alkyl, for example (C₁-C₄)-alkyl such as methyl or ethyl, optionally substituted phenyl, phenyl-(C₁-C₄)-alkyl, for example benzyl, optionally substituted in the phenyl group, hydroxy-(C₂-C₄)-alkyl such as, for example 2-hydroxyethyl, acetyl or another acyl group, methylsulfonyl or another sulfonyl group, aminocarbonyl, (C₁-C₄)-alkyloxycarbonyl, etc. In general, in the compounds of the formula I nitrogen heterocycles can also be present as N-oxides or as quaternary salts. Ring sulfur atoms can be oxidized to the sulfoxide or to the sulfone. Thus, for example a tetrahydrothienyl residue may be present as S,S-dioxotetrahydrothienyl residue or a thiomorpholinyl residue like thiomorpholin-4-yl may be present as 1-oxo-thiomorpholin-4-yl or 1,1-dioxo-thiomorpholin-4-yl. A substituted group Het that can be present in a specific position of the compounds of formula I can independently of other groups Het be substituted by substituents selected from any desired subgroup of the substituents listed before and/or in the definition of that group.
- Halogen is fluorine, chlorine, bromine or iodine, preferably fluorine, chlorine or bromine, particularly preferably chlorine or bromine.

Optically active carbon atoms present in the compounds of the formula I can independently of each other have R configuration or S configuration. The compounds of

the formula I can be present in the form of pure enantiomers or pure diastereomers or in the form of mixtures of enantiomers and/or diastereomers, for example in the form of racemates. The present invention relates to pure enantiomers and mixtures of enantiomers as well as to pure diastereomers and mixtures of diastereomers. The invention comprises mixtures of two or of more than two stereoisomers of the formula I, and it comprises all ratios of the stereoisomers in the mixtures. In case the compounds of the formula I can be present as E isomers or Z isomers (or cis isomers or trans isomers) the invention relates both to pure E isomers and pure Z isomers and to E/Z mixtures in all ratios. The invention also comprises all tautomeric forms of the compounds of the formula I.

Diastereomers, including E/Z isomers, can be separated into the individual isomers, for example, by chromatography. Racemates can be separated into the two enantiomers by customary methods, for example by chromatography on chiral phases or by resolution, for example by crystallization of diastereomeric salts obtained with optically active acids or bases. Stereochemically uniform compounds of the formula I can also be obtained by employing stereochemically uniform starting materials or by using stereoselective reactions.

The choice of incorporating into a compound of the formula I a building block with R configuration or S configuration, or in the case of an amino acid unit present in a compound of the formula I of incorporating a building block designated as D-amino acid or L-amino acid, can depend, for example, on the desired characteristics of the compound of the formula I. For example, the incorporation of a D-amino acid building block can confer increased stability in vitro or in vivo. The incorporation of a D-amino acid building block also can achieve a desired increase or decrease in the pharmacological activity of the compound. In some cases it can be desirable to allow the compound to remain active for only a short period of time. In such cases, the incorporation of an L-amino acid building block in the compound can allow endogenous peptidases in an individual to digest the compound in vivo, thereby limiting the individual's exposure to the active compound. A similar effect may also be observed in the compounds of the invention by changing the configuration in another building block from S configuration to R configuration or vice versa. By taking into consideration the

medical needs one skilled in the art can determine the desirable characteristics, for example a favorable stereochemistry, of the required compound of the invention.

Physiologically tolerable salts of the compounds of formula I are nontoxic salts that are
5 physiologically acceptable, in particular pharmaceutically utilizable salts. Such salts of compounds of the formula I containing acidic groups, for example a carboxy group COOH, are for example alkali metal salts or alkaline earth metal salts such as sodium salts, potassium salts, magnesium salts and calcium salts, and also salts with
10 physiologically tolerable quarternary ammonium ions such as tetramethylammonium or tetraethylammonium, and acid addition salts with ammonia and physiologically tolerable organic amines, such as methylamine, dimethylamine, trimethylamine, ethylamine, triethylamine, ethanolamine or tris-(2-hydroxyethyl)amine. Basic groups contained in the compounds of the formula I, for example amino groups or guanidino groups, form acid
15 addition salts, for example with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid or phosphoric acid, or with organic carboxylic acids and sulfonic acids such as formic acid, acetic acid, oxalic acid, citric acid, lactic acid, malic acid, succinic acid, malonic acid, benzoic acid, maleic acid, fumaric acid, tartaric acid, methanesulfonic acid or p-toluenesulfonic acid. Compounds of the formula I which
20 simultaneously contain a basic group and an acidic group, for example a guanidino group and a carboxy group, can also be present as zwitterions (betaines) which are likewise included in the present invention.

Salts of compounds of the formula I can be obtained by customary methods known to those skilled in the art, for example by combining a compound of the formula I with an
25 inorganic or organic acid or base in a solvent or dispersant, or from other salts by cation exchange or anion exchange. The present invention also includes all salts of the compounds of the formula I which, because of low physiological tolerability, are not directly suitable for use in pharmaceuticals but are suitable, for example, as
30 intermediates for carrying out further chemical modifications of the compounds of the formula I or as starting materials for the preparation of physiologically tolerable salts. The present invention furthermore includes all solvates of compounds of the formula I, for example hydrates or adducts with alcohols.


The invention also includes derivatives and modifications of the compounds of the formula I, for example prodrugs, protected forms and other physiologically tolerable derivatives, as well as active metabolites of the compounds of the formula I. The invention relates in particular to prodrugs and protected forms of the compounds of the formula I which can be converted into compounds of the formula I under physiological conditions. Suitable prodrugs for the compounds of the formula I, i. e. chemically modified derivatives of the compounds of the formula I having properties which are improved in a desired manner, for example with respect to solubility, bioavailability or duration of action, are known to those skilled in the art. More detailed information relating to prodrugs is found in standard literature like, for example, Design of Prodrugs, H. Bundgaard (ed.), Elsevier, 1985, , Fleisher et al., Advanced Drug Delivery Reviews 19 (1996) 115-130; or H. Bundgaard, Drugs of the Future 16 (1991) 443 which are all incorporated herein by reference. Suitable prodrugs for the compounds of the formula I are especially acyl prodrugs and carbamate prodrugs of acylatable nitrogen-containing groups such as amino groups and the guanidino group and also ester prodrugs and amide prodrugs of carboxylic acid groups which may be present in compounds of the formula I. In the acyl prodrugs and carbamate prodrugs one or more, for example one or two, hydrogen atoms on nitrogen atoms in such groups are replaced with an acyl group or a carbamate, preferably a (C₁-C₆)-alkyloxycarbonyl group. Suitable acyl groups and carbamate groups for acyl prodrugs and carbamate prodrugs are, for example, the groups R^{p1}-CO- and R^{p2}O-CO-, in which R^{p1} is hydrogen, (C₁-C₁₈)-alkyl, (C₃-C₈)-cycloalkyl, (C₃-C₈)-cycloalkyl-(C₁-C₄)-alkyl-, (C₆-C₁₄)-aryl, Het-, (C₆-C₁₄)-aryl-(C₁-C₄)-alkyl- or Het-(C₁-C₄)-alkyl- and in which R^{p2} has the meanings indicated for R^{p1} with the exception of hydrogen.

A further embodiment of the present invention are prodrugs of the compounds of the formula I, preferably (C₁-C₆)-acyl prodrugs and (C₁-C₆)-alkyloxycarbonyl prodrugs.


The present invention also relates to processes of preparation by which the compounds of the formula I are obtainable and which comprise carrying out one or more of the synthesis steps described below. The compounds of the formula I can generally be prepared, for example in the course of a convergent synthesis, by linkage of two or more fragments which can be derived retrosynthetically from the formula I. In the course of the preparation of the compounds of the formula I it can generally be advantageous or

necessary to introduce functional groups which could lead to undesired reactions or side reactions in the respective synthesis step, in the form of precursor groups which are later converted into the desired functional groups, or to temporarily block functional groups by a protective group strategy suited to the synthesis problem. Such strategies are well known to those skilled in the art (see, for example, Greene and Wuts, Protective Groups in Organic Synthesis, Wiley, 1991). As examples of precursor groups nitro groups and cyano groups may be mentioned which can later be converted by reduction, for example by catalytic hydrogenation, into amino groups and aminomethyl groups, respectively. Protective groups can also have the meaning of a solid phase, and cleavage from the solid phase stands for the removal of the protective group. The use of such techniques is known to those skilled in the art (Burgess K (Ed.) Solid Phase Organic Synthesis, New York: Wiley, 2000). For example, a phenolic hydroxy group can be attached to a trityl-polystyrene resin, which serves as a protecting group, and the molecule is cleaved from this resin by treatment with TFA at a later stage of the synthesis.

For example, for the preparation of a compound of the formula I a building block of the formula XI,



in which R_0 , Q, Q', X, are as defined above for the compounds of the formula I but functional groups can optionally also be present in the form of precursor groups or can be protected by protective groups known to those skilled in the art, e.g. an amino group can be protected with a tert.-butoxycarbonyl group or a benzyloxycarbonyl group. $R^{1'}$, $R^{1''}$, $R^{1'''}$, $R^{1''''}$, are defined as hydrogen or as R^1 which has the same meaning as in formula I but can optionally also be present in the form of precursor groups or can be protected by protective groups known to those skilled in the art, e.g. a hydroxy group may be attached to a polystyrene resin, and Y is a nucleophilically substitutable leaving group or a hydroxyl group, is reacted with a fragment of the formula III



in which R₁₀ and V are as defined above for the compounds of the formula I but functional groups can optionally also be present in the form of precursor groups or can be protected by protective groups.

- 5 The group COY in the formula XI is preferably the carboxylic acid group COOH or an activated carboxylic acid derivative. Y can thus be, for example, hydroxyl, halogen, in particular chlorine or bromine, alkoxy, in particular methoxy or ethoxy, aryloxy, for example phenoxy or pentafluorophenoxy, phenylthio, methylthio, 2-pyridylthio or a residue of a nitrogen heterocycle bonded via a nitrogen atom, in particular a residue of
10 an azole, such as, for example, 1-imidazolyl. Y can furthermore be, for example, ((C₁-C₄)-alkyl)-O-CO-O- or tolylsulfonyloxy and the activated acid derivative can thus be a mixed anhydride.

- If Y is hydroxyl, then the carboxylic acid is expediently first activated, for example by one
15 of the various methods used for peptide couplings which are well known to those skilled in the art. Examples of suitable activation agents are O-((cyano(ethoxycarbonyl)methylene)amino)-1,1,3,3-tetramethyluronium tetrafluoroborate (TOTU) ; (König et al., Proc. 21st Europ. Peptide Symp. 1990 (eds. Giralt, Andreu), Escom, Leiden 1991, p. 143), O-(benzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HBTU), O-
20 (7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HATU) (L. A. Carpino, J. Am. Chem. Soc. 1993, 115, 4397), or carbodiimides like dicyclohexylcarbodiimide or diisopropylcarbodiimide. The activation of the carboxylic acid function may also favorably be carried, for example, by conversion of the carboxylic acid group into the pentafluorophenyl ester using dicyclohexylcarbodiimide and pentafluorophenol. A
25 number of suitable methods for the preparation of activated carboxylic acid derivatives are also indicated with details of source literature in J. March, Advanced Organic Chemistry, Fourth Edition, John Wiley & Sons, 1992. The activation and the subsequent reaction with the compound of the formula III are usually carried in the presence of an inert solvent or diluent, for example DCM, chloroform, THF, diethyl ether, n-heptane, n-
30 hexane, n-pentane, cyclohexane, diisopropyl ether, methyl tert-butyl ether, acetonitrile, DMF, DMSO, dioxane, toluene, benzene, ethyl acetate or a mixture of these solvents, if appropriate with addition of a base such as, for example, potassium tert-butoxide or tributylamine or triethylamine or diisopropylethylamine.

The resulting product is a compound of the formula I in which functional groups can also be present in the form of precursor groups or can be protected by protective groups. If still any protective groups or precursor groups are present they are then removed by known methods (see Greene and Wuts, Protective Groups in Organic Synthesis, Wiley, 1991), or converted in the desired final groups, respectively. E.g. , if a carboxylic acid group protected as tert-butyl ester and the free carboxylic acid is to be prepared as the final compound the protective group can be removed by reaction with trifluoroacetic acid or tert.-butoxycarbonyl protecting groups can be removed by treatment with trifluoroacetic acid. If desired, with the obtained compounds further reactions can then be carried out by standard processes, for example acylation reactions or esterification reactions, or the compounds can be converted into physiologically tolerable salts or prodrugs by standard processes known to those skilled in the art.

Other compounds of the formula I can be prepared in a similar fashion as described above by coupling of a fragment of the formula XIII with fragment XII.



in which R_0 , Q, Q', X and Y are as defined above for the compounds of the formula I, W is the substructure of formula III, but functional groups can optionally also be present in the form of precursor groups or can be protected by protective groups known to those skilled in the art, e.g. an amino group can be protected with a tert.-butoxycarbonyl group or a benzyloxycarbonyl group or a hydroxy group may be attached to a polystyrene resin.

The fragments of the formula XI, XII and XIII are prepared by methods well known to those skilled in the art (E.g. in J March, Advanced Organic Chemistry, 4th Edition, John Wiley & Sons, 1992; RC Larock, Comprehensive Organic Transformations, VCH Publishers, New York 1989).

The compounds of the present invention are serine protease inhibitors which inhibit the activity of the blood coagulation enzymes factor Xa and/or factor VIIa. In particular, they are highly active inhibitors of factor Xa. They are specific serine protease inhibitors inasmuch as they do not substantially inhibit the activity of other proteases whose inhibition is not desired. The activity of the compounds of the formula I can be determined, for example, in the assays described below or in other assays known to

those skilled in the art. With respect to factor Xa inhibition, a preferred embodiment of the invention comprises compounds which have a $K_i \leq 1$ for factor Xa inhibition as determined in the assay described below, with or without concomitant factor VIIa inhibition, and which preferably do not substantially inhibit the activity of other proteases involved in coagulation and fibrinolysis whose inhibition is not desired (using the same concentration of the inhibitor). The compounds of the invention inhibit factor Xa catalytic activity either directly, within the prothrombinase complex or as a soluble subunit, or indirectly, by inhibiting the assembly of factor Xa into the prothrombinase complex.

- 10 The present invention also relates to the compounds of the formula I and/or their physiologically tolerable salts and/or their prodrugs for use as pharmaceuticals (or medicaments), to the use of the compounds of the formula I and/or their physiologically tolerable salts and/or their prodrugs for the production of pharmaceuticals for inhibition of factor Xa and/or factor VIIa or for influencing blood coagulation, inflammatory response
- 15 or fibrinolysis or for the therapy or prophylaxis of the diseases mentioned above or below, for example for the production of pharmaceuticals for the therapy and prophylaxis of cardiovascular disorders, thromboembolic diseases or restenoses. The invention also relates to the use of the compounds of the formula I and/or their physiologically tolerable salts and/or their prodrugs for the inhibition of factor Xa and/or factor VIIa or for
- 20 influencing blood coagulation or fibrinolysis or for the therapy or prophylaxis of the diseases mentioned above or below, for example for use in the therapy and prophylaxis of cardiovascular disorders, thromboembolic diseases or restenoses, and to methods of treatment aiming at such purposes including methods for said therapies and prophylaxis. The present invention also relates to pharmaceutical preparations (or pharmaceutical
- 25 compositions) which contain an effective amount of at least one compound of the formula I and/or its physiologically tolerable salts and/or its prodrugs in addition to a customary pharmaceutically acceptable carrier, i. e. one or more pharmaceutically acceptable carrier substances or excipients and/or auxiliary substances or additives.
- 30 Preferred are the treatment of disease states such as abnormal thrombus formation, acute myocardial infarction, unstable angina, thromboembolism, acute vessel closure associated with thrombolytic therapy or percutaneous transluminal coronary angioplasty, transient ischemic attacks, stroke, pathologic thrombus formation occurring in the veins of the lower extremities following abdominal, knee and hip surgery, a risk of pulmonary

thromboembolism, or disseminated systemic intravascular coagulopathy occurring in vascular systems during septic shock, certain viral infections or cancer.

The compounds of the formula I and their physiologically tolerable salts and their
5 prodrugs can be administered to animals, preferably to mammals, and in particular to humans as pharmaceuticals for therapy or prophylaxis. They can be administered on their own, or in mixtures with one another or in the form of pharmaceutical preparations which permit enteral or parenteral administration.

10 The pharmaceuticals can be administered orally, for example in the form of pills, tablets, lacquered tablets, coated tablets, granules, hard and soft gelatin capsules, solutions, syrups, emulsions, suspensions or aerosol mixtures. Administration, however, can also be carried out rectally, for example in the form of suppositories, or parenterally, for example intravenously, intramuscularly or subcutaneously, in the form of injection
15 solutions or infusion solutions, microcapsules, implants or rods, or percutaneously or topically, for example in the form of ointments, solutions or tinctures, or in other ways, for example in the form of aerosols or nasal sprays.

The pharmaceutical preparations according to the invention are prepared in a manner
20 known per se and familiar to one skilled in the art, pharmaceutically acceptable inert inorganic and/or organic carriers being used in addition to the compound(s) of the formula I and/or its (their) physiologically tolerable salts and/or its (their) prodrugs. For the production of pills, tablets, coated tablets and hard gelatin capsules it is possible to use, for example, lactose, corn starch or derivatives thereof, talc, stearic acid or its salts,
25 etc. Carriers for soft gelatin capsules and suppositories are, for example, fats, waxes, semisolid and liquid polyols, natural or hardened oils, etc. Suitable carriers for the production of solutions, for example injection solutions, or of emulsions or syrups are, for example, water, saline, alcohols, glycerol, polyols, sucrose, invert sugar, glucose, vegetable oils, etc. Suitable carriers for microcapsules, implants or rods are, for example,
30 copolymers of glycolic acid and lactic acid. The pharmaceutical preparations normally contain about 0.5 % to 90 % by weight of the compounds of the formula I and/or their physiologically tolerable salts and/or their prodrugs. The amount of the active ingredient of the formula I and/or its physiologically tolerable salts and/or its prodrugs in the

pharmaceutical preparations normally is from about 0.5 mg to about 1000 mg, preferably from about 1 mg to about 500 mg.

In addition to the active ingredients of the formula I and/or their physiologically acceptable salts and/or prodrugs and to carrier substances, the pharmaceutical preparations can contain additives such as, for example, fillers, disintegrants, binders, lubricants, wetting agents, stabilizers, emulsifiers, preservatives, sweeteners, colorants, flavorings, aromatizers, thickeners, diluents, buffer substances, solvents, solubilizers, agents for achieving a depot effect, salts for altering the osmotic pressure, coating agents or antioxidants. They can also contain two or more compounds of the formula I and/or their physiologically tolerable salts and/or their prodrugs. In case a pharmaceutical preparation contains two or more compounds of the formula I the selection of the individual compounds can aim at a specific overall pharmacological profile of the pharmaceutical preparation. For example, a highly potent compound with a shorter duration of action may be combined with a long-acting compound of lower potency. The flexibility permitted with respect to the choice of substituents in the compounds of the formula I allows a great deal of control over the biological and physico-chemical properties of the compounds and thus allows the selection of such desired compounds. Furthermore, in addition to at least one compound of the formula I and/or its physiologically tolerable salts and/or its prodrugs, the pharmaceutical preparations can also contain one or more other therapeutically or prophylactically active ingredients.

As inhibitors of factor Xa and/or factor VIIa the compounds of the formula I and their physiologically tolerable salts and their prodrugs are generally suitable for the therapy and prophylaxis of conditions in which the activity of factor Xa and/or factor VIIa plays a role or has an undesired extent, or which can favorably be influenced by inhibiting factor Xa and/or factor VIIa or decreasing their activities, or for the prevention, alleviation or cure of which an inhibition of factor Xa and/or factor VIIa or a decrease in their activity is desired by the physician. As inhibition of factor Xa and/or factor VIIa influences blood coagulation and fibrinolysis, the compounds of the formula I and their physiologically tolerable salts and their prodrugs are generally suitable for reducing blood clotting, or for the therapy and prophylaxis of conditions in which the activity of the blood coagulation system plays a role or has an undesired extent, or which can favorably be influenced by reducing blood clotting, or for the prevention, alleviation or cure of which a decreased

activity of the blood coagulation system is desired by the physician. A specific subject of the present invention thus are the reduction or inhibition of unwanted blood clotting, in particular in an individual, by administering an effective amount of a compound I or a physiologically tolerable salt or a prodrug thereof, as well as pharmaceutical preparations therefor.

Conditions in which a compound of the formula I can be favorably used include, for example, cardiovascular disorders, thromboembolic diseases or complications associated, for example, with infection or surgery. The compounds of the present invention can also be used to reduce an inflammatory response. Examples of specific disorders for the treatment or prophylaxis of which the compounds of the formula I can be used are coronary heart disease, myocardial infarction, angina pectoris, vascular restenosis, for example restenosis following angioplasty like PTCA, adult respiratory distress syndrome, multi-organ failure, stroke and disseminated intravascular clotting disorder. Examples of related complications associated with surgery are thromboses like deep vein and proximal vein thrombosis which can occur following surgery. In view of their pharmacological activity the compounds of the invention can replace or supplement other anticoagulant agents such as heparin. The use of a compound of the invention can result, for example, in a cost saving as compared to other anticoagulants.

When using the compounds of the formula I the dose can vary within wide limits and, as is customary and is known to the physician, is to be suited to the individual conditions in each individual case. It depends, for example, on the specific compound employed, on the nature and severity of the disease to be treated, on the mode and the schedule of administration, or on whether an acute or chronic condition is treated or whether prophylaxis is carried out. An appropriate dosage can be established using clinical approaches well known in the medical art. In general, the daily dose for achieving the desired results in an adult weighing about 75 kg is from 0.01 mg/kg to 100 mg/kg, preferably from 0.1 mg/kg to 50 mg/kg, in particular from 0.1 mg/kg to 10 mg/kg, (in each case in mg per kg of body weight). The daily dose can be divided, in particular in the case of the administration of relatively large amounts, into several, for example 2, 3 or 4, part administrations. As usual, depending on individual behavior it may be necessary to deviate upwards or downwards from the daily dose indicated.

A compound of the formula I can also advantageously be used as an anticoagulant outside an individual. For example, an effective amount of a compound of the invention can be contacted with a freshly drawn blood sample to prevent coagulation of the blood sample. Further, a compound of the formula I and its salts can be used for diagnostic purposes, for example in in vitro diagnoses, and as an auxiliary in biochemical investigations. For example, a compound of the formula I can be used in an assay to identify the presence of factor Xa and/or factor VIIa or to isolate factor Xa and/or factor VIIa in a substantially purified form. A compound of the invention can be labeled with, for example, a radioisotope, and the labeled compound bound to factor Xa and/or factor VIIa is then detected using a routine method useful for detecting the particular label. Thus, a compound of the formula I or a salt thereof can be used as a probe to detect the location or amount of factor Xa and/or factor VIIa activity in vivo, in vitro or ex vivo.

Furthermore, the compounds of the formula I can be used as synthesis intermediates for the preparation of other compounds, in particular of other pharmaceutical active ingredients, which are obtainable from the compounds of the formula I, for example by introduction of substituents or modification of functional groups.

It is understood that changes that do not substantially affect the activity of the various embodiments of this invention are included within the invention disclosed herein. Thus, the following examples are intended to illustrate but not limit the present invention.

Examples


Abbreviations used:

25	tert-Butyl	tBu
	Dichloromethane	DCM
	Diethyl azodicarboxylate	DEAD
	Diisopropyl azodicarboxylate	DIAD
	N,N'-Diisopropylcarbodiimide	DIC
30	N,N-Diisopropyl-N-ethylamine	DIEA
	N,N-Dimethylformamide	DMF
	Dimethylsulfoxide	DMSO
	O-(7-Azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium-hexafluorophosphate	HATU

1-Hydroxy-7-azabenzotriazole	HOAt
N-Ethylmorpholine	NEM
Methanol	MeOH
Tetrahydrofuran	THF
5 Trifluoroacetic acid	TFA

When in the final step of the synthesis of a compound an acid such as trifluoroacetic acid or acetic acid was used, for example when trifluoroacetic acid was employed to remove a tert-butyl group or when a compound was purified by chromatography using an eluent which contained such an acid, in some cases, depending on the work-up procedure, for example the details of a freeze-drying process, the compound was obtained partially or completely in the form of a salt of the acid used, for example in the form of the acetic acid salt or trifluoroacetic acid salt or hydrochloric acid salt.

15 Example 1: (S)-(1-Carbamimidoyl-piperidin-4-yl)-{3-[2-(2,4-dichloro-phenyl)-ethoxy]-5-hydroxy- benzoylamino}-acetic acid methyl ester



(a) 3,5-Dihydroxy-benzoic acid allyl ester

To 1.5 g of 3,5-dihydroxybenzoic acid in screw-capped vial was added 10 g of allyl alcohol and the vial was closed and cooled to -20°C . To the cold contents of the reaction vial was added 5 mL of trimethylsilyl chloride via syringe through a septum. The reaction vial was allowed to warm to room temperature and agitated for 16 hours. The vial was opened carefully and its contents transferred to a round-bottomed flask. The solvent was removed under reduced pressure and the residual solid was dried under reduced pressure over potassium hydroxide pellets for 12 h. The semisolid product was used in the subsequent synthetic steps without further purification. The product was analyzed by HPLC and had a retention time of 3.65 on 5 cm C_{18} reversed phase column with flow rate of 2.5 mL/min of pure acetonitrile (for solvent A) and 0.1 % aqueous trifluoroacetic acid (for solvent B). The product was characterized by ^1H NMR (DMSO-d_6 ,

350 MHz): δ = 6.87 (s, 2H, aromatic); 6.44 (s, 1H, aromatic); 5.85-5.96 (m, 1H); 5.16-5.33 (m, 2H); 4.65-4.67 (dd, 2H).

(b) 300 mg of (2'-chloro)-Chlorotriptyl-polystyrene resin (0.39 mmols; loading 1.3 mmol/g Cl) was treated with 4 mL of dichloroethane and the resin was left to swell at room temperature for 30 minutes. The solvent was removed by filtration and the resin was treated with a solution of 227 mg of 3,5-dihydroxy-benzoic acid allyl ester and 0.4 mL of DIEA in 5 mL anhydrous dichloromethane. The resin suspension was agitated for 3 to 4 h at 60° C. The resin was washed with DMF (3 times), DCM (5 times) and DMF (5 times) and used in next step.

(c) The resin from step (b) was washed with anhydrous THF (3 times) and suspended in 4 mL of anhydrous THF containing 511 mg of triphenylphosphine and 745 mg of 2,4-dichlorophenethyl alcohol. The suspension was cooled to -15 °C and 0.384 mL of DIAD was added. The resin suspension was agitated at room temperature for 12 h. The solvent was removed by filtration and the resin was washed with THF (9 times), DMF (5 times), DCM (5 times). The resin was used in the next step.

(d) The resin from step (c) was suspended in DCM and 365 mg of 1,3-dimethylbarbituric acid was added in the presence of 45 mg of Pd (0)(PPh₃)₄ under argon. The resin suspension was agitated for one hour at room temperature. The solvent was removed by filtration and the resin was washed with DCM and dried.


(e) Dried resin from step (d) was washed with DMF and suspended in 3 mL DMF containing 265 mg of HOAt and 0.302 mL of DIC. The resin was agitated for 5 minutes and 613 mg of (S)-amino-[1-(tert-butoxycarbonylamino-imino-methyl)-piperidin-4-yl]-acetic acid methyl ester was added. The resin suspension was agitated for 12 h. The resin was washed with DMF and DCM and dried under reduced pressure for 6 - 8 h.

(f) The dried resin from step (e) was suspended in DCM containing 50% TFA and agitated at room temperature for 45 minutes. The resin suspension was filtered, washed with DCM: TFA (1:1) and the washings combined with the cleavage filtrate. The cleavage solution was dried under reduced pressure. The solid product was lyophilized from 30% aqueous acetonitrile and crude product purified by HPLC on reverse phase C₁₈ column.

Fractions containing the desired product were pooled and lyophilized to give (S)-(1-Carbamimidoyl-piperidin-4-yl)-{3-[2-(2,4-dichloro-phenyl)-ethoxy]-5-hydroxy-benzoylamino}-acetic acid methyl ester as a white solid. The product was identified by LC/MS to give $m/e = 523 (M+H)^+$.

5

Example 2: 4-Bromo-N-(1-carbamimidoyl-piperidin-4-ylmethyl)-3-[2-(2,4-dichloro-phenyl)-ethoxy]-5- hydroxy-benzamide



(a) 4-Bromo-3,5-dihydroxy-benzoic acid allyl ester

- 10 This compound was prepared analogously to 3,5-Dihydroxy-benzoic acid allyl ester [Example 1(a)], however 4-bromo-3,5-dihydroxybenzoic acid was used instead of 3,5-dihydroxybenzoic acid. ^1H NMR (DMSO- d_6 , 350 MHz): $\delta = 7.04$ (s, 2H, aromatic); 5.97-6.06 (m, 1H); 5.25-5.40 (m, 2H); 4.73-4.75(dd, 2H).


- 15 (b) The title compound was synthesized analogously to Example 1, steps (b)-(f) with the following differences:

In step (b) 4-Bromo-3,5-dihydroxy-benzoic acid allyl ester was used instead of 3,5-dihydroxy-benzoic acid allyl ester;

- 20 In step (e) [(4-aminomethyl-piperidin-1-yl)-imino-methyl]-carbamic acid tert-butyl ester was used instead of (S)-amino-[1-(tert-butoxycarbonylamino-imino-methyl)-piperidin-4-yl]-acetic acid methyl ester. The final product was purified by HPLC and characterized by LC/MS to give $m/e=543.3 (M+H)^+$.

Example 3: N-(1-Carbamimidoyl-piperidin-4-ylmethyl)-3-[2-(2,4-dichloro-phenyl)-ethoxy]-5-hydroxy-4- methyl-benzamide

25



(a) 3,5-dihydroxy-4-methylbenzoic acid allyl ester






This compound was prepared analogously to 3,5-Dihydroxy-benzoic acid allyl ester [Example 1(a)], however 3,5-dihydroxy-4-methylbenzoic acid was used instead of 3,5-dihydroxybenzoic acid.

(b) The title compound was synthesized analogously to Example 2, with the following difference:


3,5-dihydroxy-4-methyl-benzoic acid allyl ester was used instead of 4-Bromo-3,5-dihydroxy-benzoic acid allyl ester. The final product was purified by HPLC and characterized by LC/MS to give $m/e=478.8$ ($M+H$)⁺.

Analogously to the above examples the following example compounds were prepared by similar procedures and characterized by LC/MS:

Example	Structure	MWt	(M+H) ⁺
4		464.13	465.3
5		536.15	537.3

6		564.10	565.3
7		513.22	514.3
8		523.10	524
9		411.23	412
10		475.11	476

Example 11: N-(1-Carbamimidoyl-piperidin-4-ylmethyl)-3-[2-(2,4-dichloro-phenyl)-ethoxy]-4-methoxy- benzamide



(a) 3-Hydroxy-4-methoxy-benzoic acid methyl ester

- 5 10 mL of thionyl chloride was added to 250 mL of methanol at 0 °C. The solution was stirred for 10 minutes and 25 g of 3-hydroxy-4-methoxybenzoic acid were added. The reaction was stirred for 16 h at room temperature then heated to 50 °C for 3 h. The solvents were removed under reduced pressure. The residue was used directly in the next step.

10

(b) 3-[2-(2,4-Dichlorophenyl)-ethoxy]-4-methoxy-benzoic acid methyl ester


- 20g of triphenylphosphine and 10 g of 3-Hydroxy-4-methoxy-benzoic acid methyl ester were dissolved in 200 mL of anhydrous THF. The solution was cooled to 0 °C to 10 °C and a solution of 11.4mL DEAD in 30 mL anhydrous THF was added dropwise over 20
15 min. The reaction was warmed to room temperature and stirred for 45 min. A solution of 11.3 mL 2-(2,4-Dichlorophenyl)-ethanol in 10mL anhydrous THF was added with cooling. The reaction was stirred at room temperature for 16 h, then the solvents were removed under reduced pressure. The residue was treated with n-heptane:ethyl acetate/1:1. The filtrate was dried under reduced pressure. The product was purified by silica gel
20 chromatography, eluting with n-heptane:ethyl acetate/4:1, then n-heptane:ethyl acetate/3:1. Yield 17g.

(c) 3-[2-(2,4-Dichlorophenyl)-ethoxy]-4-methoxy-benzoic acid

- 17 g of 3-[2-(2,4-Dichlorophenyl)-ethoxy]-4-methoxy-benzoic acid methyl ester was
25 dissolved in 200 mL of methanol:water/3:1. 4.1 g of lithium hydroxide monohydrate was added to the solution, and the reaction was stirred at room temperature for 16 h then at 90 °C for 2 h. The solution was cooled to room temperature, then acidified with half-concentrated hydrochloric acid. The solvents were removed under reduced pressure and the residue was washed twice with warm water to remove salts.


- (d) A solution of 100 mg 3-[2-(2,4-Dichlorophenyl)-ethoxy]-4-methoxy-benzoic acid in 2 ml DMF was activated by the addition of 53 mg carbonyldiimidazole. After stirring for 2 h at RT 90 mg of 4-Aminomethyl-piperidine-1-carboxamide hydrochloride and 2 ml DMSO were added and the mixture was stirred overnight. Subsequent dilution with 3 ml water and filtration through a chem elut® cartridge, eluting with ethyl acetate yielded after concentration under reduced pressure a white solid. Purification by preparative HPLC (C₁₈ reverse phase column, elution with a H₂O/MeCN gradient with 0.5 % TFA) and lyophilisation afforded 30 mg (20%) of the title compound as a white powder.
- MS (ESI +) m/e 479.3 (M+H) chloro pattern.

Example 12: 3-[2-(2,4-Dichloro-phenyl)-ethoxy]-N-[1-(1-imino-ethyl)-piperidin-4-ylmethyl]-4-methyl- benzamide; compound with trifluoro-acetic acid



- 50 mg of 3-[2-(2,4-Dichloro-phenyl)-ethoxy]-4-methyl-benzoic acid and 58.9 mg of C-[1-(1-imino-ethyl)-piperidin-4-yl]-methylamine di-trifluoroacetic acid salt were dissolved in 5 ml of DMF. After cooling to 0 °C 64.3 mg of HATU and 70.8 mg of NEM were added. The mixture was stirred at 0°C for 2 hours and at room temperature for 3 hours. The solvent was removed under reduced pressure and the residue was purified by preparative HPLC (C18 reverse phase column) (gradient acetonitrile water (containing 0.1% trifluoro-acetic acid) 90:10 to 0:100). The fractions containing the product were evaporated and lyophilized. Yield: 48 mg (54%), MS: 462.2/464.3 (M+H)⁺.


Example 13: N-(1-Carbamimidoyl-piperidin-4-ylmethyl)-3-[2-(2,4-dichloro-phenyl)-ethoxy]-4-methyl- benzamide; compound with trifluoro-acetic acid



50 mg of 3-[2-(2,4-Dichloro-phenyl)-ethoxy]-4-methyl-benzoic acid and 35.2 mg of 4-aminomethyl-piperidine-1-carboxamide dihydrochloride were dissolved in 5 ml of DMF. After cooling to 0 °C 64.3 mg of HATU and 70.8 mg of NEM were added. The mixture was stirred at 0 °C for 2 hours and at room temperature for 3 hours. The solvent was removed under reduced pressure and the residue was purified by preparative HPLC (C₁₈ reverse phase column) (gradient acetonitrile water (containing 0.1% trifluoro-acetic acid) 90:10 to 0:100). The fractions containing the product were evaporated and lyophilized.

Yield: 45 mg (51%), MS: 463.3/465.3 (M+H)⁺.

Example 14: {Amino-[4-({3-[2-(2,4-dichloro-phenyl)-ethoxy]-4-methyl-benzoylamino}-methyl)-piperidin-1-yl]-methylene}-carbamic acid benzyl ester



97 mg of 3-[2-(2,4-Dichloro-phenyl)-ethoxy]-4-methyl-benzoic acid and 124.3 mg of [Amino-(4-aminomethyl-piperidin-1-yl)-methylene]-carbamic acid benzyl ester trifluoroacetic acid salt were dissolved in 5 ml of DMF. After cooling to 0 °C 136 mg of HATU and 106 mg of NEM were added. The mixture was stirred at 0 °C for 2 hours and at room temperature for 3 hours. The solvent was removed under reduced pressure and the residue was purified by preparative HPLC (C₁₈ reverse phase column) (gradient acetonitrile water (containing 0.1% trifluoro-acetic acid) 90:10 to 0:100). The fractions containing the product were evaporated and lyophilized.

Yield: 146 mg (79.5%), MS: 597.3/599.3 (M+H)⁺.

Pharmacological testing

5

The ability of the compounds of the formula I to inhibit factor Xa or factor VIIa or other enzymes like thrombin, plasmin, or trypsin can be assessed by determining the concentration of the compound of the formula I that inhibits enzyme activity by 50 %, i. e. the IC₅₀ value, which is related to the inhibition constant Ki. Purified enzymes are used in
10 chromogenic assays. The concentration of inhibitor that causes a 50 % decrease in the rate of substrate hydrolysis is determined by linear regression after plotting the relative rates of hydrolysis (compared to the uninhibited control) versus the log of the concentration of the compound of formula I. For calculating the inhibition constant Ki, the IC₅₀ value is corrected for competition with substrate using the formula

15

$$K_i = IC_{50} / \{1 + (\text{substrate concentration} / K_m)\}$$

wherein Km is the Michaelis-Menten constant (Chen and Prusoff, Biochem. Pharmacol. 22 (1973), 3099-3108; I. H. Segal, Enzyme Kinetics, 1975, John Wiley & Sons, New York, 100-125; which are incorporated herein by reference).

20

a) Factor Xa Assay

In the assay for determining the inhibition of factor Xa activity TBS-PEG buffer (50 mM Tris-HCl, pH 7.8, 200 mM NaCl, 0.05 % (w/v) PEG-8000, 0.02 % (w/v) NaN₃) was used. The IC₅₀ was determined by combining in appropriate wells of a Costar half-area microtiter plate 25 µl human factor Xa (Enzyme Research Laboratories, Inc.; South
25 Bend, Indiana) in TBS-PEG; 40 µl 10 % (v/v) DMSO in TBS-PEG (uninhibited control) or various concentrations of the compound to be tested diluted in 10 % (v/v) DMSO in TBS-PEG; and substrate S-2765 (N(α)-benzyloxycarbonyl-D-Arg-Gly-L-Arg-p-nitroanilide; Kabi Pharmacia, Inc.; Franklin, Ohio) in TBS-PEG.

30

The assay was performed by pre-incubating the compound of formula I plus enzyme for 10 min. Then the assay was initiated by adding substrate to obtain a final volume of 100 µl. The initial velocity of chromogenic substrate hydrolysis was measured by the change in absorbance at 405 nm using a Bio-tek Instruments kinetic plate reader (Ceres UV900HDI) at 25 °C during the linear portion of the time course (usually 1.5 min after

addition of substrate). The enzyme concentration was 0.5 nM and substrate concentration was 140 μ M.

b) Factor VIIa Assay


- 5 The inhibitory activity towards factor VIIa/tissue factor activity was determined using a chromogenic assay essentially as described previously (J. A. Ostrem et al., Biochemistry 37 (1998) 1053-1059 which is incorporated herein by reference). Kinetic assays were conducted at 25 °C in half-area microtiter plates (Costar Corp., Cambridge, Massachusetts) using a kinetic plate reader (Molecular Devices Spectramax 250). A
- 10 typical assay consisted of 25 μ l human factor VIIa and TF (5 nM and 10 nM, respective final concentration) combined with 40 μ l of inhibitor dilutions in 10% DMSO/TBS-PEG buffer (50 mM Tris, 15 mM NaCl, 5 mM CaCl_2 , 0.05 % PEG 8000, pH 8.15). Following a 15 minute preincubation period, the assay was initiated by the addition of 35 μ l of the chromogenic substrate S-2288 (D-Ile-Pro-Arg-p-nitroanilide, Pharmacia Hepar Inc., 500
- 15 μ M final concentration).

The following test results (inhibition constants K_i (FXa) for inhibition of factor Xa) were obtained:

- 20 Example 1: K_i (FXa) 2.116 micromolar
Example 2: K_i (FXa) 0.0137 micromolar
Example 3: K_i (FXa) 0.0585 micromolar
Example 4: K_i (FXa) 0.4635 micromolar
Example 6: K_i (FXa) 2.280 micromolar
- 25 Example 11: K_i (FXa) 0.189 micromolar
Example 12: K_i (FXa) 0.14 micromolar
Example 13: K_i (FXa) 0.173 micromolar
Example 14: K_i (FXa) 5.077 micromolar (Prodrug)

Patent Claims

1. A compound of the formula I,




wherein

- 5 R_0 is 1. phenyl, wherein phenyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R^2 or
2. a mono- or bicyclic 5- to 10-membered heteroaryl containing one or two nitrogen atoms as ring heteroatoms, wherein heteroaryl is unsubstituted or mono-, di- or trisubstituted independently of one
- 10 another by R^2 ,
- R^2 is 1. $-\text{NO}_2$,
2. halogen,
3. $-\text{CN}$,
4. $-\text{OH}$,
- 15 5. $-\text{NH}_2$,
6. $(\text{C}_1\text{-C}_8)\text{-alkyloxy-}$, wherein alkyloxy is unsubstituted or mono-, di- or trisubstituted independently of one another by halogen, amino group, hydroxy group or methoxy group, or
7. $(\text{C}_1\text{-C}_8)\text{-alkyl-}$, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by halogen, amino
- 20 group, hydroxy group or methoxy group,
- Q and Q' are independently of one another identical or different and are a direct bond, $-\text{O-}$, $-\text{S-}$, $-\text{NR}^{10}\text{-}$, $-\text{C(O)NR}^{10}\text{-}$, $-\text{NR}^{10}\text{C(O)-}$, $-\text{S(O)-}$, $-\text{SO}_2\text{-}$, $-\text{NR}^{10}\text{-SO}_2\text{-}$, $-\text{SO}_2\text{-NR}^{10}\text{-}$ oder $-\text{C(O)-}$;
- 25 R^{10} is hydrogen atom or $(\text{C}_1\text{-C}_4)\text{-alkyl-}$,
- X is 1. a direct bond,
2. $(\text{C}_1\text{-C}_6)\text{-alkylen-}$, wherein alkylen is unsubstituted or mono-, di- or trisubstituted independently of one another by halogen, amino group or a hydroxy group,

3. (C₃-C₆)-cycloalkylen, wherein cycloalkylen is unsubstituted or mono-, di- or trisubstituted independently of one another by halogen, amino group or a hydroxy group,

provided that at least one of Q, X and Q' is not a direct bond,

- 5 D is an atom out of the group carbon, oxygen, sulfur and nitrogen,
the substructure of formula III



(III)

- is 1. a mono- or bicyclic 5- to 10-membered carbocyclic aryl group, wherein said 5- to 10-membered carbocyclic aryl group is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹,
- 10 2. phenyl, wherein phenyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹,
3. a mono- or bicyclic 5-to 10-membered heterocyclic group (Het), containing one or more heteroatoms as ring heteroatoms, such as nitrogen, sulfur or oxygen, wherein said Het group is unsubstituted or mono-, di- or
- 15 trisubstituted independently of one another by R¹, or
4. pyridyl, wherein pyridyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹,

- R¹ is 1. halogen,
2. -NO₂,
- 20 3. -CN,
4. R¹¹R¹²N-, wherein R¹¹R¹² independently of one another are hydrogen atom, (C₁-C₄)-alkyl- or (C₁-C₆)-acyl-,
5. (C₁-C₈)-alkylamino-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³,
- 25 6. -OH,
7. -SO₂-NH₂,
8. (C₁-C₈)-alkyloxy-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³,
9. (C₆-C₁₄)-aryl, wherein aryl is unsubstituted or mono-, di- or
- 30 trisubstituted independently of one another by R¹³,

10. (C₁-C₈)-alkyl-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³,
11. hydroxycarbonyl-(C₁-C₈)-alkylureido-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³,
- 5 12. (C₁-C₈)-alkyloxycarbonyl-(C₁-C₈)-alkylureido-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³,
13. (C₁-C₈)-alkylsulfonyl-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³, or
- 10 14. -C(O)-NR¹⁴R¹⁵, wherein R¹⁴R¹⁵ independently of one another are hydrogen atom or (C₁-C₄)-alkyl-, or


two R¹ residues bonded to adjacent ring carbon atoms together with the carbon atoms to which they are bonded form an aromatic ring condensed to the ring depicted in formula I, where the ring formed by the two R¹ residues is

15 unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³, R¹¹ and R¹² together with the nitrogen atom to which they are bonded form a saturated or unsaturated 5- to 6-membered monocyclic heterocyclic ring which in addition to the nitrogen atom carrying R¹¹ and R¹² can contain one or two identical or different ring heteroatoms chosen from oxygen, sulfur and nitrogen, and in


20 which one or two of the ring carbon atoms can be substituted by oxo to form -C(O)- residue(s),

- R¹³ is
1. halogen,
 2. -NO₂,
 3. -CN,
 - 25 4. -OH,
 5. (C₁-C₈)-alkyl-,
 6. (C₁-C₈)-alkyloxy-,
 7. -CF₃ or
 8. -NH₂,


30 V is a residue of the formulae IIa, IIb, IIc, IId, IIe or II f,




IIa




IIb




IIe



IIc



IIId



IIIf

wherein

L is a direct bond or (C₁-C₃)-alkylene, wherein alkylene is unsubstituted or mono-, di- or trisubstituted independently of one another by A,

- A is
1. hydrogen atom,
 2. -C(O)-OH,
 3. -C(O)-O-(C₁-C₄)-alkyl, wherein alkyl is unsubstituted or mono-, di- or tri- substituted independently of one another by -OH, -NH₂ or -(C₁-C₄)-alkoxy,
 4. -C(O)-NR⁴R⁵,
 5. (C₁-C₄)-alkyl-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by -OH, -NH₂ or -(C₁-C₄)-alkoxy,
 6. -SO₂-NH₂ or
 7. -SO₂-CH₃,

U is -NH₂, (C₁-C₄)-alkyl-, -NH-C(O)-O-(C₁-C₄)-alkyl or -NH-C(O)-O-(C₁-C₄)-alkyl-aryl,

M is hydrogen atom, (C₁-C₃)-alkyl- or -OH,

R⁴ and R⁵ are independently of one another identical or different and are

1. hydrogen atom,
2. (C₁-C₁₂)-alkyl-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³ as defined above,

3. (C₆-C₁₄)-aryl-(C₁-C₄)-alkyl-, wherein alkyl and aryl are unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³ as defined above,
 4. (C₆-C₁₄)-aryl-, wherein aryl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³ as defined above,
 5. Het-, wherein Het- is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³ as defined above, or
 6. Het-(C₁-C₄)-alkyl-, wherein alkyl and Het- are unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³ as defined above, or
- 10 R⁴ and R⁵ together with the nitrogen atom to which they are bonded form a saturated 3- to 8-membered monocyclic heterocyclic ring which in addition to the nitrogen atom carrying R⁴ and R⁵ can contain one or two identical or different ring heteroatoms chosen from oxygen, sulfur and nitrogen;
- 15 in all its stereoisomeric forms and mixtures thereof in any ratio, and its physiologically tolerable salts.

2. A compound of formula I as claimed in claim 1, wherein

R₀ is phenyl, wherein phenyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R², or
 20 pyridyl, wherein pyridyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R²,

- R² is
1. -NO₂,
 2. halogen,
 - 25 3. -CN,
 4. -OH,
 5. -NH₂,
 6. (C₁-C₄)-alkyloxy-, wherein alkyloxy is unsubstituted or mono-, di- or trisubstituted independently of one another by halogen, amino group, hydroxy group or methoxy group, or
 - 30 7. -(C₁-C₄)-alkyl, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by halogen, amino group, hydroxy group or methoxy group,

Q, Q', X, R¹, R¹¹ and R¹² are as defined in claim 1,

D is an atom out of the group carbon and nitrogen,
the substructure of formula III is

1. phenyl, wherein phenyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R^1 , or
2. pyridyl, wherein pyridyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R^1 ,

R^{13} is

1. halogen,
2. $-NO_2$,
3. $-CN$,
4. $-OH$,
5. (C_1-C_4) -alkyl-,
6. (C_1-C_4) -alkyloxy-,
7. $-CF_3$ or
8. $-NH_2$,

R_{10} is hydrogen atom or methyl,

V is a fragment of the formula IIa, IIb, IIc, IId, IIe or II f as defined above, wherein L, U, M, R^4 and R^5 are as defined in claim 1, and

A is

1. hydrogen atom,
2. $-C(O)-OH$,
3. $-C(O)-O-(C_1-C_4)$ -alkyl, wherein alkyl is unsubstituted or mono-, di- or tri- substituted independently of one another by $-OH$, $-NH_2$ or $-(C_1-C_4)$ -alkoxy,
4. $-C(O)-NR^4R^5$ or
5. (C_1-C_4) -alkyl-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by $-OH$, $-NH_2$ or $-(C_1-C_4)$ -alkoxy.

3. A compound of the formula I as claimed in claims 1 or 2, wherein

R^0 is phenyl, wherein phenyl is mono-, di- or trisubstituted independently of one another by R^2 , or
pyridyl, wherein pyridyl is mono-, di- or trisubstituted independently of one another by R^2 ,

R^2 is

1. $-NH_2$,

2. halogen,
3. -CN,
4. -OH,
5. (C₁-C₄)-alkyloxy-, wherein alkyloxy is unsubstituted or substituted by an amino group, or
6. -(C₁-C₄)-alkyl, wherein alkyl is unsubstituted or substituted by an amino group,

Q and Q' are independently of one another identical or different and are a direct bond, -O-, -C(O)NR¹⁰-, -NR¹⁰C(O)-; -NR¹⁰-SO₂-; or -SO₂-NR¹⁰-;

- X is
1. a direct bond or
 2. (C₁-C₄)-alkylen, wherein alkylen is unsubstituted or mono-, di- or tri-substituted independently of one another by halogen, amino group or a hydroxy group,

D is an atom out of the group carbon and nitrogen,

the substructure of formula III is

phenyl or pyridyl, wherein phenyl and pyridyl are unsubstituted or mono-, di- or trisubstituted independently of one another by R¹,

- R¹ is
1. halogen,
 2. -NO₂,
 3. -CN,
 4. -NH₂,
 5. (C₁-C₄)-alkylamino-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³,
 6. -OH,
 7. -SO₂-NH₂,
 8. (C₁-C₄)-alkyloxy-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³,
 9. (C₆-C₁₄)-aryl, wherein aryl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³,
 10. (C₁-C₄)-alkyl-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³,
 11. (C₁-C₄)-alkylsulfonyl-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³,

12. $-C(O)-NR^{14}R^{15}$, wherein $R^{14}R^{15}$ independently of one another are hydrogen atom or (C_1-C_4) -alkyl-,

13. $R^{11}R^{12}N-$, wherein R^{11} and R^{12} are as defined above, or

14. $-NR^4R^5$,

5

R^{13} is 1. halogen,

2. $-NO_2$,

3. $-CN$,

4. $-OH$,

5. (C_1-C_4) -alkyl-,

10

6. (C_1-C_4) -alkyloxy-,

7. $-CF_3$ or

8. $-NH_2$,

R_{10} is hydrogen atom or methyl,

V is a fragment of the formula IIa, IIb, IIc, IId, IIe or II f as defined above, wherein

15

L is a direct bond or (C_1-C_3) -alkylen-,

A is hydrogen atom, $-C(O)-OH$, $-C(O)-O-(C_1-C_4)$ -alkyl, $-C(O)-NR^4R^5$ or (C_1-C_4) -alkyl-,

U is $-NH_2$, methyl, $-NH-C(O)-O-(C_1-C_4)$ -alkyl or $-NH-C(O)-O-(CH_2)$ -phenyl,

M is hydrogen atom, (C_1-C_3) -alkyl- or $-OH$, and

20

R^4 and R^5 are independently of one another hydrogen atom or (C_1-C_4) -alkyl-.

4. A compound of formula I as claimed in one or more of claims 1 to 3, wherein

25

R_0 is phenyl or pyridyl, wherein phenyl and pyridyl independently from one another are mono-, di- or trisubstituted independently of one another by R^2 ,

R^2 is 1. halogen,

2. $-CN$,

3. (C_1-C_4) -alkyloxy-, wherein alkyloxy is unsubstituted or substituted by halogen or an amino group, or

30

4. $-(C_1-C_4)$ -alkyl, wherein alkyl is unsubstituted or substituted by an amino group or halogen,

Q and Q' are independently of one another identical or different and are a direct bond, $-O-$, $-C(O)NR^{10}$ -, $-NR^{10}C(O)-$; $-NR^{10}-SO_2-$; or $-SO_2-NR^{10}-$;

X is $-(C_1-C_3)$ -alkylen-, wherein alkylen is unsubstituted or mono-, di- or trisubstituted independently of one another by halogen, amino group or hydroxy group,

D is the atom carbon,

the substructure of formula III is

phenyl, wherein phenyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R^1 ,

R^1 is 1. halogen,

2. $-NO_2$,

3. $-CN$,

4. $-NH_2$,

5. (C_1-C_4) -alkylamino-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R^{13} ,

6. $-OH$,

7. $-SO_2-NH_2$,

8. (C_1-C_4) -alkyloxy-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R^{13} ,

9. (C_6-C_{14}) -aryl, wherein aryl is unsubstituted or mono-, di- or trisubstituted independently of one another by R^{13} ,

10. (C_1-C_4) -alkyl-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R^{13} ,

11. (C_1-C_4) -alkylsulfonyl-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R^{13} ,

12. $-C(O)-NR^{14}R^{15}$, wherein $R^{14}R^{15}$ independently of one another are hydrogen atom or (C_1-C_4) -alkyl-,

13. $R^{11}R^{12}N-$, wherein R^{11} and R^{12} are as defined above, or

14. $-NR^4R^5$,

R^{13} is 1. halogen,

2. $-CF_3$,

3. $-NH_2$,

4. $-OH$,

5. (C_1-C_4) -alkyl- or

6. (C_1-C_4) -alkyloxy-,

R_{10} is hydrogen atom, and

V is a fragment of the formula IIa, IIb, IIc or IId as defined above, wherein

L is a direct bond or (C₁-C₂)-alkylen-,

A is hydrogen atom, -C(O)-OH, -C(O)-O-(C₁-C₄)-alkyl, -C(O)-NR⁴R⁵ or (C₁-C₄)-alkyl,

5 U is -NH₂, methyl, -NH-C(O)-O-(C₁-C₄)-alkyl or -NH-C(O)-O-(CH₂)-phenyl,

M is hydrogen atom or (C₁-C₃)-alkyl-, and

R⁴ and R⁵ are independently of one another hydrogen atom or methyl.

10 5. A compound of formula I as claimed in one or more of claims 1 to 4, wherein

R₀ is phenyl, wherein phenyl is mono-, di- or trisubstituted independently of one another by R²,

R² is 1. halogen,

2. (C₁-C₄)-alkyloxy-, wherein alkyloxy is unsubstituted or substituted by halogen or an amino group, or

15 3. -(C₁-C₄)-alkyl, wherein alkyl is unsubstituted or substituted by an amino group or halogen,

Q and Q' are independently of one another identical or different and are a direct bond, -O-, -C(O)NR¹⁰-, -NR¹⁰C(O)-; -NR¹⁰-SO₂-; or -SO₂-NR¹⁰-;

20 X is -(C₁-C₃)-alkylen-,

D is the atom carbon,

the substructure of formula III is

phenyl, wherein phenyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹,

25 R¹ is 1. halogen,

2. -NO₂,

3. -CN,

4. -NH₂,

5. (C₁-C₄)-alkylamino-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³,

30

6. -OH,

7. -SO₂-NH₂,

8. (C₁-C₄)-alkyloxy-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³,

9. (C₁-C₄)-alkyl-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³,
10. (C₁-C₄)-alkylsulfonyl-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R¹³,
11. -C(O)-NR¹⁴R¹⁵, wherein R¹⁴R¹⁵ independently of one another are hydrogen atom or (C₁-C₂)-alkyl-,
12. R¹¹R¹²N-, wherein R¹¹ and R¹² are as defined above, or
13. -NR⁴R⁵,

- R¹³ is
1. halogen,
 2. -CF₃,
 3. -NH₂,
 4. -OH,
 5. (C₁-C₄)-alkyl- or
 6. (C₁-C₄)-alkyloxy-,

R₁₀ is hydrogen atom, and

V is a fragment of the formula IIa, IIb, IIc or IId as defined above, wherein

L is a direct bond or (C₁-C₂)-alkylen-,

A is hydrogen atom, -C(O)-OH, -C(O)-O-(C₁-C₄)-alkyl, -C(O)-NR⁴R⁵ or -(C₁-C₄)-alkyl,

U is -NH₂, methyl, -NH-C(O)-O-(C₁-C₄)-alkyl or -NH-C(O)-O-(CH₂)-phenyl,

M is hydrogen atom or methyl, and

R⁴ and R⁵ are independently of one another hydrogen atom or methyl.

6. A compound of formula I as claimed in one or more of claims 1 to 5, wherein R₀ is phenyl, wherein phenyl is disubstituted independently of one another by R²,

- R² is
1. halogen,
 2. (C₁-C₂)-alkyloxy-, wherein alkyloxy is unsubstituted or substituted by an amino group, or
 3. -(C₁-C₄)-alkyl, wherein alkyl is unsubstituted or substituted by an amino group,

Q and Q' are independently of one another identical or different and are a direct bond or -O-,

X is $-\text{CH}_2-\text{CH}_2-$,

D is the atom carbon,

the substructure of formula III is

phenyl, wherein phenyl is unsubstituted or mono-, di- or trisubstituted

independently of one another by R^1 ,

R^1 is 1. halogen,

2. $-\text{OH}$,

3. $-\text{NH}_2$,

4. $-\text{C}(\text{O})-\text{NR}^{14}\text{R}^{15}$, wherein $\text{R}^{14}\text{R}^{15}$ independently of one another are hydrogen atom or (C_1-C_2) -alkyl-,

5. (C_1-C_3) -alkyloxy-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R^{13} , or

6. (C_1-C_3) -alkyl-, wherein alkyl is unsubstituted or mono-, di- or trisubstituted independently of one another by R^{13} ,

R^{13} is fluorine or chlorine,

R_{10} is hydrogen atom, and

V is a fragment of the formula IIa, IIb, IIc or IId as defined above, wherein

L is a direct bond or (C_1-C_2) -alkylen-,

A is hydrogen atom, $-\text{C}(\text{O})-\text{OH}$, $-\text{C}(\text{O})-\text{O}-(\text{C}_1-\text{C}_4)$ -alkyl, $-\text{C}(\text{O})-\text{NR}^4\text{R}^5$ or $-(\text{C}_1-\text{C}_4)$ -alkyl,


U is $-\text{NH}_2$, methyl, $-\text{NH}-\text{C}(\text{O})-\text{O}-(\text{C}_1-\text{C}_4)$ -alkyl or $-\text{NH}-\text{C}(\text{O})-\text{O}-(\text{CH}_2)$ -phenyl,

M is hydrogen atom, and

R^4 and R^5 are independently of one another hydrogen atom or methyl.

7. A process for the preparation of a compound of the formula I as claimed in one or more of claims 1 to 6, comprises

a) linking a building block of the formula XI,




in which R_0 , Q, Q' and X, are as in claims 1 to 6, and $R^{1'}$, $R^{1''}$, $R^{1'''}$, $R^{1''''}$, are hydrogen atom or as R^1 as defined in claims 1 to 6, but where in R_0 , Q, R^1 Q' and X functional groups can also be present in protected form or in the form of precursor groups, and Y is a nucleophilically substituable leaving group or a hydroxyl group,

is reacted with a fragment of the formula III



in which R_{10} and V are as defined in claims 1 to 6, but where in R_{10} and V functional groups can also be present in protected form or in the form of precursor groups, or

b) by coupling of a fragment of the formula XIII with fragment XII,



in which R_0 , Q, Q' and X, are as in claims 1 to 6, W is the substructure of formula III, but where in R_0 , Q, Q', W and X functional groups can also be present in protected form or in the form of precursor groups, and Y is a nucleophilically substituable leaving group or a hydroxyl group or a hydroxy group may be attached to a polystyrene resin.

8. A pharmaceutical preparation, comprising at least one compound of the formula I as claimed in one or more of claims 1 to 6 and/or its physiologically tolerable salts and a pharmaceutically acceptable carrier.
9. The use of a compound of the formula I as claimed in one or more of claims 1 to 6 and/or their physiologically tolerable salts and/or their prodrugs for the production of pharmaceuticals for inhibition of factor Xa and/or factor VIIa or for influencing blood coagulation or fibrinolysis.
10. The use as claimed in claim 9 for influencing blood coagulation, inflammatory response, fibrinolysis, cardiovascular disorders, thromboembolic diseases, restenoses, abnormal thrombus formation, acute myocardial infarction, unstable angina, acute vessel closure associated with thrombolytic therapy, thromboembolism, percutaneous, pathologic thrombus formation occurring in the veins of the lower extremities following abdominal, knee and hip surgery,

transluminal coronary angioplasty, transient ischemic attacks, stroke a risk of pulmonary thromboembolism, certain viral infections or cancer, intravascular coagulopathy occurring in vascular systems during septic shock, coronary heart disease, myocardial infarction, angina pectoris, vascular restenosis, for example restenosis following angioplasty like PTCA, adult respiratory distress syndrome, multi-organ failure, stroke and disseminated intravascular clotting disorder, thromboses like deep vein and proximal vein thrombosis which can occur following surgery.

- 5
- 10 11. A prodrug of the compound of the formula I as claimed in claims 1 to 6, preferably a (C₁-C₆)-acyl or (C₁-C₆)-alkyloxycarbonyl prodrugs of the compound of the formula I as claimed in claims 1 to 6.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/13874

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D211/34 C07D401/12 A61K31/445

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, BEILSTEIN Data, WPI Data, PAJ, EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 96 04267 A (MENARINI LAB ;MAULEON CASELLAS DAVID (ES); CARGANICO GERMANO (ES);) 15 February 1996 (1996-02-15) the whole document ----	1-11
A	WO 95 29189 A (SELECTIDE CORP) 2 November 1995 (1995-11-02) cited in the application the whole document ----	1-11
X	WO 99 48870 A (KLEIN SCOTT I ;GUERTIN KEVIN R (US); RHONE POULENC RORER PHARMA (U) 30 September 1999 (1999-09-30) see examples 12, and 3rd and 4th compounds on page 61 and claims ----- -/--	1-3,8-10

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

° Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

9 April 2002

Date of mailing of the international search report

16/04/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Scruton-Evans, I

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/13874

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 01 19788 A (COR THERAPEUTICS INC) 22 March 2001 (2001-03-22) see especially claim 3 definitions ----	1-11
A	WO 97 28129 A (PRESTON JOHN ;STOCKER ANDREW (GB); ZENECA LTD (GB); SMITHERS MICHA) 7 August 1997 (1997-08-07) the whole document -----	1-11
A	US 5 792 769 A (BONE ROGER F ET AL) 11 August 1998 (1998-08-11) the whole document -----	1-11

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1-10 (partly), 11

Present claims 1-10 relate to an extremely large number of possible compounds, due to all of the variables for Q, X and Q'. Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compounds claimed, that is those wherein Q-X-Q' is (CH₂)₂₋₃-O or CH(NH₂)-CH₂-O and where D is 1,3-phenylene. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely those parts relating to the compounds wherein Q'-W-Q is O-(CH)₁₋₆ and D is 1,3-phenylene. Furthermore, the claim 11 relates to a compound defined by reference to a desirable result or property, namely a prodrug of a compound of formula I. The claims cover all compounds having this characteristic or property, whereas the application provides support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for only a very limited number of such compounds. i.e. compound 14. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the product/compound/method/apparatus by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/13874

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9604267	A	15-02-1996	ES 2103181 A1	16-08-1997
			AT 201678 T	15-06-2001
			AU 3253995 A	04-03-1996
			DE 69521132 D1	05-07-2001
			DE 69521132 T2	27-09-2001
			DK 775133 T3	30-07-2001
			WO 9604267 A1	15-02-1996
			EP 0775133 A1	28-05-1997
			ES 2156944 T3	01-08-2001
			PT 775133 T	30-10-2001
WO 9529189	A	02-11-1995	AU 707653 B2	15-07-1999
			AU 2368395 A	16-11-1995
			CA 2186497 A1	02-11-1995
			CN 1147261 A	09-04-1997
			CZ 9603140 A3	14-05-1997
			EE 9600146 A	15-04-1997
			EP 0758341 A1	19-02-1997
			FI 964317 A	25-10-1996
			HU 76346 A2	28-08-1997
			IL 113505 A	31-12-1999
			JP 10503477 T	31-03-1998
			LT 96151 A , B	26-05-1997
			LV 11740 A	20-04-1997
			LV 11740 B	20-12-1997
			NO 964553 A	27-12-1996
			NZ 284977 A	24-09-1998
			PL 317067 A1	03-03-1997
			RU 2152954 C1	20-07-2000
			SI 9520044 A	31-10-1997
			SK 136696 A3	07-05-1997
			TW 409129 B	21-10-2000
			WO 9529189 A1	02-11-1995
			US 5849510 A	15-12-1998
			ZA 9503361 A	12-01-1996
WO 9948870	A	30-09-1999	AU 3109499 A	18-10-1999
			BR 9909086 A	04-09-2001
			CA 2325471 A1	30-09-1999
			CN 1297440 T	30-05-2001
			EP 1080075 A1	07-03-2001
			HU 0101482 A2	28-09-2001
			PL 344056 A1	24-09-2001
			TR 200002740 T2	21-12-2000
			WO 9948870 A1	30-09-1999
			US 6277865 B1	21-08-2001
			US 2002016339 A1	07-02-2002
WO 0119788	A	22-03-2001	AU 7486600 A	17-04-2001
			AU 7486700 A	17-04-2001
			WO 0119798 A2	22-03-2001
			WO 0119788 A2	22-03-2001
			AU 4535301 A	12-09-2001
			AU 5078301 A	12-09-2001
			WO 0164642 A2	07-09-2001
			WO 0164643 A2	07-09-2001
			US 2002002183 A1	03-01-2002

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/13874

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9728129	A	07-08-1997	AU 1608597 A	22-08-1997
			EP 0880502 A1	02-12-1998
			WO 9728129 A1	07-08-1997
			JP 2000504337 T	11-04-2000
			ZA 9700697 A	04-08-1997
<hr/>				
US 5792769	A	11-08-1998	AU 714292 B2	23-12-1999
			AU 7201296 A	17-04-1997
			BR 9610876 A	13-07-1999
			CA 2233344 A1	03-04-1997
			CZ 9800922 A3	15-07-1998
			EP 0859607 A1	26-08-1998
			HU 9802764 A2	29-03-1999
			JP 11512722 T	02-11-1999
			NO 981393 A	25-05-1998
			PL 325954 A1	17-08-1998
			WO 9711693 A1	03-04-1997
<hr/>				